

# Effects of *Hypericum perforatum* extract on 6-hydroxydopamine neurotoxicity in differentiated SH-SY5Y cells

Baris Bitmez<sup>a</sup>, Seda K. Gultekin<sup>a</sup>, Irem G. Albayrak<sup>a</sup>, Yigit Deveci<sup>b</sup>, Yusuf Sicak<sup>c</sup>, Emine Akalin<sup>d</sup>, Adami F. Pirhan<sup>e</sup>, Ulas Gurer<sup>a</sup>, Belkis A. Arslan<sup>a</sup>

<sup>a</sup>Department of Molecular Biology and Genetics, Uskudar University, <sup>d</sup>Department of Pharmaceutical Botany, Istanbul University, Istanbul, <sup>b</sup>Department of Chemistry, <sup>c</sup>Department of Medicinal and Aromatic Plants, Mugla Sıtkı Kocman University, Mugla, <sup>e</sup>Department of Biology, Ege University, İzmir, Turkey

Correspondence to Belkis A. Arslan, PhD, Department of Molecular Biology and Genetics, Uskudar University, Istanbul 34662, Turkey. Tel: +90 216 400 2222; fax: 90 216 474 1256; e-mail: belkisatasever.arslan@uskudar.edu.tr

**Received:** 6 December 2022

**Revised:** 13 January 2023

**Accepted:** 13 January 2023

**Published:** 27 June 2023

**Egyptian Pharmaceutical Journal** 2023, 22:188–191

## Background and objective

Parkinson's disease (PD) is the second most common neurodegenerative disease after Alzheimer's disease. In our study, PD model was created as a result of exposure to 6-hydroxydopamine (6-OHDA) in SH-SY5Y cells, which is a human neuroblastoma cell line. The protective effect of *Hypericum perforatum* on PD was investigated.

## Materials and methods

Phytochemical analysis of *H. perforatum* extract was performed. Then, SH-SY5Y cells were differentiated using retinoic acid and then administered 6-OHDA neurotoxin. To determine the protective effects of *H. perforatum* extract, we investigated the changes in the mRNA expression level of caspase-3, total oxidant status, and antioxidant levels in differentiated SH-SY5Y.

## Results and conclusion

According to our results, *H. perforatum* extract contains glycosides, tannins, flavonoids, and carbohydrates as the major secondary metabolites. *H. perforatum* extract significantly reduced caspase-3 gene expression against 6-OHDA toxicity in differentiated SH-SY5Y cells. It was found that total oxidant status level increased significantly in the 6-OHDA experimental group compared with the control and *H. perforatum* experimental groups.

It was found that *H. perforatum* extract has an inhibitory effect on caspase-3 gene expression, which plays an important role in apoptosis. Therefore, *H. perforatum* extract has been shown to have a therapeutic potential against 6-OHDA toxicity.

## Keywords:

6-hydroxydopamine, *Hypericum perforatum*, Parkinson's, disease, retinoic acid, SH-SY5Y cell line

Egypt Pharmaceut J 22:188–191

© 2023 Egyptian Pharmaceutical Journal  
1687-4315

## Introduction

Parkinson's disease (PD) is a neurodegenerative disorder characterized by the formation of Lewy bodies, which are composed of aggregated alpha-synuclein protein [1,2].

To study neurotoxicity and evaluate the effectiveness of therapeutic agents, a common practice is to use 6-hydroxydopamine (6-OHDA), a hydroxylated analog of dopamine, as a toxin in PD models [3,4].

*Hypericum perforatum* plant has been found to contain various pharmacologically and biologically active compounds [5–7]. Studies have suggested that *H. perforatum* extract may have therapeutic potential in the treatment of PD [8,9].

This study aimed to investigate changes in the mRNA expression levels of caspase-3, as well as total oxidant and antioxidant levels, in SH-SY5Y cells differentiated with retinoic acid and treated with 6-OHDA neurotoxin.

## Materials and methods

### Preparation of *Hypericum perforatum* extract

*H. perforatum* plants that were collected from Ege University's Bornova Campus in the Izmir Province of Turkey have been identified by Asst. Prof. Dr. A. Fahri Pirhan from the Ege University Faculty of Science, Department of Biology. The flower parts of the plant were dried at room temperature before being extracted using a Soxhlet apparatus with methanol. The methanol extraction was performed as described in a previous study by Deveci *et al.* [10].

### Phytochemical screening

The presence of phytochemicals was analyzed in the *H. perforatum* extract. For the screening of alkaloids, saponins, tannins, glycosides, phenols, carbohydrates,

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

proteins, and flavonoids, the method established by Khuda *et al.* [11] was used.

#### Differentiation of SH-SY5Y human neuroblastoma cells

SH-SY5Y cells were obtained from American Type Culture Collection (ATCC). The cell passage was performed according to the method described in a previous study by Kalkan *et al.* [12].

To differentiate 75% confluent SH-SY5Y cells, after 24 h culturing the cells, the ratio of fetal bovine serum in the DMEM culture medium was reduced from 10 to 5% and differentiation was initiated by adding 10  $\mu$ M retinoic acid (EMD Biosciences cat# 554720, Darmstadt, Germany). On the fourth day, the fetal bovine serum rate was reduced to 2.5%, and then on the 7th day, it was reduced to 1%. After 3 days, the experimental groups were formed [12].

#### Preparation of experimental groups

On 10th day of differentiation of SH-SY5Y cells, only the medium was placed in the first group as control. To induce the *in vitro* model of PD as the second group, 6-OHDA (Sigma H116) was adjusted to a final concentration of 100  $\mu$ M [13].

*H. perforatum* extract was added at a dose of 20  $\mu$ g/ml to the third group and after 24 h of incubation, 6-OHDA was adjusted to a final concentration of 100  $\mu$ M. To the fourth group, at a dose of 20  $\mu$ g/ml *H. perforatum* extract was added [14].

#### RT-PCR experiments

RNA isolation from the experimental groups was performed using the Zymo Quick RNA Miniprep Kit (Cat no: R1054). At the end of the isolation process, 30  $\mu$ l of nuclease-free water was added, and the RNAs were stored at  $-80^{\circ}\text{C}$ . To determine the expression levels of the caspase-3 gene for each group, the GoTaq 1-Step RT-qPCR System Technical Manual kit was used. The glyceraldehyde-3-phosphate dehydrogenase gene was used as a reference gene for normalization [15].

#### Total antioxidant status measurement

Total antioxidant status levels of experimental groups' supernatants were measured using a Rel Assay Diagnostics kit (Cat no: RL0017), according to the manufacturer's instruction. The color intensities of the samples were measured at 660-nm absorbance. Trolox, which is a vitamin E analog, was used as a stable antioxidant standard solution. The results obtained were expressed in  $\mu$ mol/l.

$$\text{Results} = \frac{[\Delta\text{Abs Sample}]}{[\Delta\text{Abs Standard}]} \times 10^*$$

(\* = Concentration of standard).

#### Total oxidant status measurement

Total oxidant status level measurements of the samples' supernatants were made with a Rel Assay Diagnostics kit (Cat no: RL0024), according to the manufacturer's instruction. The color intensities of the samples were measured at 530-nm absorbance. The assay was calibrated using a hydrogen peroxide standard solution. The results obtained were expressed in  $\mu$ mol/l.

$$\text{Results} = \frac{[\Delta\text{Abs H}_2\text{O} - n\text{Abs Sample}]}{[\Delta\text{abs H}_2\text{O} - n\text{Abs Standard}]}$$

#### Statistical analysis

Data were expressed as mean  $\pm$  SD. Statistical analysis of the results was performed using the Statistical Package for the Social Sciences (SPSS, Newyork, USA) program. Statistically significant differences were analyzed using one-way analysis of variance. The differences between the groups were determined by the least significant difference test (post-hoc test in analysis of variance). Significance was defined as a *P* value less than or equal to 0.05.

## Results

According to the results of the phytochemical screening, it was found that the *H. perforatum* extract contains glycosides, tannins, flavonoids, and carbohydrates as the major secondary metabolites (Table 1).

It was observed that the toxicity induced by 6-OHDA significantly increased the expression of the caspase-3 gene by 4.805  $\pm$  0.957-fold in the cells. However, the *H. perforatum* extract was found to significantly reduce the expression of the caspase-3 gene in the face of 6-OHDA toxicity in the differentiated SH-SY5Y

**Table 1** Phytochemical screening of the *Hypericum perforatum* extract

Phytochemicals	Chemical tests	Results
Alkaloids	Hager's test	-
Saponins	Foam test	-
Flavonoids	General test	+
Tannins	Alkaline Reagent test	+
Glycosides	Keller-Kiliani test	+
Carbohydrates	Benedict's test	+ <sup>a</sup>
Proteins	Xanthoproteic test	-

<sup>a</sup>Green (0.1–0.5% sugar).

cells. In the experimental group, a change of  $0.905 \pm 0.281$  was found in the expression of the caspase-3 gene in comparison with the control group. Furthermore, when differentiated SH-SY5Y cells were incubated only with the *H. perforatum* extract, the fold change in the expression of the caspase-3 gene was found to be  $0.704 \pm 0.049$  in comparison with the control group (Fig. 1).

In the study, it was observed that the level of total oxidant status increased significantly in the 6-OHDA experimental group in comparison with the control and *H. perforatum* experimental groups (Fig. 2). However, it has been reported that there is a decrease in the level of total antioxidant status, but no significant results were obtained. No significant changes were observed in the other groups. In the study, no significant results were obtained as a result of the application of *H. perforatum* and in the other groups (Fig. 3).

**Discussion**


The neuroprotective effects of flavonoids and tannins have been demonstrated in various studies [16,17]. Hesperidin, in particular, has been shown to have multiple beneficial pharmacological effects such as anti-oxidation, anti-inflammation, and neuroprotection [18]. The main active compounds found in the *H. perforatum* plant consist of naphthodianthrone derivatives, particularly hypericin-pseudohypericin and hyperforin, which is a phloroglucinol derivative, as well as flavonoids [7].

However, the effects of specific tannin compounds and their combinations with flavonoids on the content of *H. perforatum* are currently unknown.

In a recent study, it has been reported that increased caspase-3 activation and expression confirm the role of apoptosis in PD [19]. Additionally, another study has reported that p38 activation underlies the toxic effects of 6-OHDA on dopaminergic neurons and that activation of the p38 signaling pathway promotes


Downloaded from http://ejournals.ewp.com/egpj by BMDM5ePHKav1ZEoum1lQIN4e+kLlHEZgbsHh04XIM0hCwWCX1AW nYOp/llQH33D00dRv/7TVSf4C3VC1Y0abggQZXdgg/2MwZLel= on 08/11/2023

Figure 1




Caspase-3 gene expression alterations among experimental groups.

Figure 2



Total oxidant status levels in control of the experimental groups.

Figure 3



Total antioxidant status levels in control of the experimental groups.

cell apoptosis, a process known to be involved in PD and plays a role in neuroinflammation [20,21]. The findings obtained in these studies suggest that the content of *H. perforatum* extract has the potential as a promising new molecule for the treatment of PD.

In a study conducted by Gomez del Rio *et al.* [8], it was shown that a hyperforin-rich *H. perforatum* extract prevents dopaminergic neuron loss in experimental animals in a Parkinson's model created with rotenone. Various studies have also shown that 6-OHDA stimulation increases p38 phosphorylation and caspase-3 activation [22,23].

## Conclusion

According to the results of our study, it was found that the *H. perforatum* extract has an inhibitory effect on caspase-3 gene expression, which plays an important role in apoptosis. Therefore, it has been demonstrated that *H. perforatum* extract has therapeutic potential against 6-OHDA toxicity. However, further research is necessary to fully understand their specific and combinatorial roles in neuroprotection by isolating the tannins and flavonoids found in the extract.

## Acknowledgements

Funding: This study was supported by the Scientific and Technological Research Council of Turkey (TUBITAK) 2209-A (Project No: 1919B012000056).

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

- Wang SF, Liu LF, Wu MY, Cai CZ, Su H, Tan J, *et al.* Baicalein prevents 6-OHDA/ascorbic acid-induced calcium-dependent dopaminergic neuronal cell death. *Sci Rep* 2017; 7:8398.
- Kesh S, Kannan RR, Sivaji K, Balakrishnan A. Hesperidin downregulates kinases Irfk2 and gsk3 $\beta$  in a 6-OHDA induced Parkinson's disease model. *Neurosci Lett* 2021; 740:135426.
- Kondo M, Hara H, Kamijo F, Kamiya T, Adachi T. 6-Hydroxydopamine disrupts cellular copper homeostasis in human neuroblastoma SH-SY5Y cells. *Metallomics* 2021; 13:041.
- Xu Z, Yang D, Huang X, Huang H. Astragaloside IV protects 6-hydroxydopamine-induced SH-SY5Y cell model of Parkinson's disease via activating the JAK2/STAT3 pathway. *Front Neurosci* 2021; 15:631501.
- Bilia AR, Gallori S, Vincieri FF. St. John's wort and depression: efficacy, safety and tolerability—an update. *Life Sci* 2002; 70:3077–3096.
- Greeson JM, Sanford B, Monti DA. St. John's wort (*Hypericum perforatum*): a review of the current pharmacological, toxicological, and clinical literature. *Psychopharmacology (Berl)* 2001; 153:402–414.
- Shakya P, Marslin G, Siram K, Beerhues L, Franklin G. Elicitation as a tool to improve the profiles of high-value secondary metabolites and pharmacological properties of *Hypericum perforatum*. *J Pharma Pharmacol* 2017; 71:70–82.
- Gomez del Rio MA, Sánchez-Reus MI, Iglesias I, Pozo MA, García-Arencibia M, Fernández-Ruiz J, *et al.* Neuroprotective properties of standardized extracts of *Hypericum perforatum* on rotenone model of Parkinson's disease. *CNS Neurol Disord Drug Targets* 2013; 12:665–679.
- Vecchia DD, Schamne MG, Ferro MM, Santos AFCD, Latyki CL, Lara DVD, *et al.* Effects of *Hypericum perforatum* on turning behavior in an animal model of Parkinson's disease. *Braz J Pharma Sci* 2015; 51:111–115.
- Deveci Y, Günel Sadık G, Akalın E, Kuşoğlu Gültekin S, Yanık A, Atasever Arslan B. Anti-diabetic effects of *Berberis cretica* extract in INS-1E cells. *Int J Sci Lett* 2022; 3:121–128.
- Khuda F, Alam N, Khalil AAK, Jan A, Naureen F, Ullah Z, *et al.* Screening of *Rhamnus purpurea* (Edgew.) leaves for antimicrobial, antioxidant, and cytotoxic potential. *ACS Omega* 2022; 7:22977–22985.
- Kalkan Z, Durasi IM, Sezerman U, Atasever Arslan B. Potential of GRID2 receptor gene for preventing TNF-induced neurodegeneration in autism. *Neurosci Lett* 2016; 620:62–69.
- Wei L, Ding L, Mo MS, Lei M, Zhang L, Chen K, Xu P. Wnt3a protects SH-SY5Y cells against 6-hydroxydopamine toxicity by restoration of mitochondria function. *Transl Neurodegener* 2015; 4:11.
- Akkus RY, Bitmez B, Kusoglu Gültekin S, Albayrak IG, Ozen F, Deveci Y, *et al.* Neuroprotective effect of *Hypericum perforatum* extract against aluminum-maltolate induced toxicity in SH-SY5Y cells. *Int J Sci Lett* 2022; 4:277–291.
- Sang Q, Liu X, Wang L, Qi L, Sun W, Wang W, *et al.* CircSNCA downregulation by pramipexole treatment mediates cell apoptosis and autophagy in Parkinson's disease by targeting miR-7. *Aging* 2018; 10:1281–1293.
- Hussain G, Huang J, Rasul A, Anwar H, Imran A, Maqbool J, *et al.* roles of plant-derived tannins in neurodegenerative and neuropsychiatry disorders: an updated review. *Molecules* 2019; 24:2213.
- Potshangbam AM, Rathore RS, Nongdam P. Discovery of sulfone-resistant dihydropterolate synthase (DHPS) as a target enzyme for kaempferol, a natural flavanoid. *Heliyon* 2020; 6:e03378.
- Welbat JU, Naewla S, Pannangrong W, Sirichoat A, Aranarochana A, Wigmore P. Neuroprotective effects of hesperidin against methotrexate-induced changes in neurogenesis and oxidative stress in the adult rat. *Biochem Pharmacol* 2020; 178:114083.
- Erekat NS. Apoptosis and its role in Parkinson's disease. *Exon Publ* 2018; 4:65–82.
- Obergasteiger J, Frapporti G, Pramstaller PP, Hicks AA, Volta M. A new hypothesis for Parkinson's disease pathogenesis: GTPase-p38 MAPK signaling and autophagy as convergence points of etiology and genomics. *Mol Neurodegener* 2018; 13:1–17.
- Wang ZQ, Li K, Huang J, Huo TT, Lv PY. MicroRNA Let-7i is a promising serum biomarker for post-stroke cognitive impairment and alleviated OGD-induced cell damage in vitro by regulating Bcl-2. *Front Neurosci* 2020; 14:215.
- Jaisin Y, Ratanachamnong P, Kuanpradit C, Khumpum W, Suksamram S. Protective effects of  $\gamma$ -mangostin on 6-OHDA-induced toxicity in SH-SY5Y cells. *Neurosci Lett* 2018; 665:229–235.
- Cirmi S, Maueri A, Lombardo GE, Russo C, Musumeci L, Gangemi S, *et al.* A flavonoid-rich extract of mandarin juice counteracts 6-OHDA-induced oxidative stress in SH-SY5Y cells and modulates Parkinson-related genes. *Antioxidants* 2021; 10:539.