

## Some Growth Parameters of Five Fish Species in the Lower Sakarya River, Turkey

İsmail Reis<sup>1</sup> , Hasan Cerim<sup>1</sup> , Celal Ateş<sup>1</sup> 

Cite this article as: Reis, İ., Cerim, H., Ateş, C. (2020). Some growth parameters of five fish species in the lower Sakarya river, Turkey. *Aquatic Sciences and Engineering*, 35(3), 75-82.

### ABSTRACT

In this study, a total of 1283 samples of five fish species belonging to two families, Cyprinidae and Leuciscidae, were collected from the Lower Sakarya River between June 2017 and May 2018 in order to determine some growth parameters. The samples were collected monthly with trammel net, fykenets, and electro shocker. The age of the fish was determined from the scales. The von Bertalanffy's growth model was calculated  $L_t = 92.18(1 - e^{-0.054(t+0.040)})$  for *A. brama*,  $L_t = 69.40(1 - e^{-0.040(t+0.030)})$  for *B. bjoerkna*,  $L_t = 51.09(1 - e^{-0.114(t+0.024)})$  for *C. gibelio*,  $L_t = 48.11(1 - e^{-0.088(t+0.023)})$  for *R. rutilus* and  $L_t = 41.74(1 - e^{-0.104(t+0.035)})$  for *V. vimba*. The phi-prime growth performance index ( $\Phi'$ ) value was computed as 2.628, 2.268, 2.474, 2.307 and 2.260 for *A. brama*, *B. bjoerkna*, *C. gibelio*, *R. rutilus* and *V. vimba*, respectively. This study provides basic information on some growth parameters of five fish species living in the Lower Sakarya River. The results of this study are useful for fishery managements and stock assessment in the Sakarya River.

**Keywords:** Age and growth, cyprinidae, leuciscidae, fishery management, Sakarya River

ORCID IDs of the author:  
İ.R. 0000-0003-4599-6780;  
H.C. 0000-0003-3025-1444;  
C.A. 0000-0002-7336-0387

<sup>1</sup>Muğla Sıtkı Koçman University  
Fisheries Faculty, Department of  
Fishing Technology, Muğla, Turkey

Submitted:  
26.11.2019

Revision Requested:  
29.02.2020

Revision Requested:  
09.03.2020

Accepted:  
20.03.2020

Online published:  
06.04.2020

Correspondence:  
İsmail Reis  
E-mail:  
[ismailreis@mu.edu.tr](mailto:ismailreis@mu.edu.tr)

©Copyright 2020 by Aquatic  
Sciences and Engineering  
Available online at  
<https://dergipark.org.tr/ase>

### INTRODUCTION

Rivers and natural lakes are important ecosystems of our world and cover approximately 2.5% of the earth's surface (Shiklomanov, 1999). Turkey has important freshwater resources and one of these freshwater resources is the Sakarya River. The Sakarya River basin (58000 km<sup>2</sup>), one of the major rivers pouring into the Black Sea, covers approximately 7 % of Turkey's surface area (783000 km<sup>2</sup>). Its average flow rate is about 190 m<sup>3</sup> per second. The water temperature changes between 7 to 24 °C through out the year. The river basin is divided into three regions named Lower, Mid and Upper Basin (Şengörür & İsa, 2001).

Cyprinidae is found in North America (from northern Canada to southern Mexico), Africa, and Eurasia. Cyprinidae is the largest family of freshwater fish with 346 genera and 3,170 species in the world. Leuciscidae is the other important

freshwater fish with 90 genera and 672 species (Eschmeyer, Fricke & van der Laan, 2017). Various researches have been carried out on the fish species living in the Sakarya River and its tributaries. (Ölmez, 1992; Emiroğlu, 2011; Kahraman, Gök-türk & Aydın, 2014; Korkmaz & Zencir Tanır, 2016; Memiş, Tosun, Yamaner, Tunçelli & Gessner, 2019; Reis, Cerim & Ateş, 2019).

Age and growth are related with each other. Age gives a knowledge about sexual maturity, spawning period, fish size, growth rate and lifespan. Knowledge of all these parameters are important data for fisheries management and vary among populations. Accurate age determination and estimates of growth parameters are fundamental requirements for understanding population dynamics and maintaining sustainable yields in fisheries biology (Campana & Thorrold, 2001).

In this study, some growth parameters were determined for *Abramis brama* (Linnaeus, 1758),

*Blicca bjoerkna* (Linnaeus, 1758), *Carassius gibelio* (Bloch, 1782), *Rutilus rutilus* (Linnaeus, 1758) and *Vimba vimba* (Linnaeus, 1758) that were caught in the lower Sakarya River. These data contribute to the sustainable management of the Sakarya River fisheries.

## MATERIALS AND METHODS

This study was carried out between June 2017 and May 2018 in the 159.5 km section of the Sakarya River within the borders of Sakarya province. The aforementioned section includes Meceke in the south of Pamukova and Karasu Yenimahalle, where it deposits into the Black Sea.

The samples were collected monthly with trammel nets (inner panel: 52-72-88 mm, outer panel: 300 mm; stretched mesh sized), fyke net (140 mm stretched mesh sized, 5 m leader net) and electro shocker (SAMUS 1000; 500W) from the three stations (Pamukova, Adapazarı and Karasu) identified in the lower Sakarya River Basin (Figure 1). The sampling areas were sandy-muddy substrates and depths were between 1.5-10 meters.




Figure 1. Study area.

The samples were brought to the laboratory and the fish species were determined according to their diagnostic characteristics (Kottelat & Freyhof, 2007). Total lengths and weights of samples were measured with measuring boards (0.1 cm) and precision balance (0.01 g). Scales were used to determine the age of the fish. The scales were taken from the area between the dorsalfin

and the lateral line on the left side of the fish by forceps and placed in numbered envelopes (Lagler, 1966). Scales were removed from the envelopes and placed in petri dishes, containing 3% NaOH solution, in order to be purified from foreign bodies. Randomly selected scales were examined under a binocular microscope (Chugunova, 1963).

Growth parameters were investigated by applying the von Bertalanffy growth function. The von Bertalanffy growth function was calculated as follows:  $L_t = L_{\infty} (1 - e^{-k(t-t_0)})$  (von Bertalanffy, 1957), where  $L_t$  is length at age  $t$ ,  $L_{\infty}$  is asymptotic length,  $k$  is the growth coefficient, and  $t_0$  is the hypothetical age at which length is equal to zero (Ricker, 1975).

The growth performance index was calculated by the equation of Pauly & Munro (1984):

$$\phi' = \text{Log } k + 2 \text{ Log } L_{\infty}$$

## RESULTS AND DISCUSSION

In this study, all samples were analyzed to estimate age and growth parameters, including five fish species from the Lower Sakarya River, Turkey. The parameters shown in Table 1 included sample size ( $n$ ), range of total length (TL) and body weight ( $W$ ), and standard error (SE).

The von Bertalanffy's growth model was calculated  $L_t = 92.18(1 - e^{-0.054(t+0.040)})$  for *A. brama*,  $L_t = 69.40(1 - e^{-0.040(t+0.030)})$  for *B. bjoerkna*,  $L_t = 51.09(1 - e^{-0.114(t+0.024)})$  for *C. gibelio*,  $L_t = 48.11(1 - e^{-0.088(t+0.023)})$  for *R. rutilus* and  $L_t = 41.74(1 - e^{-0.104(t+0.035)})$  for *V. vimba*. The asymptotic length ( $L_{\infty}$ ), growth coefficient ( $k$ ), hypothetical age ( $t_0$ ) and growth performance index ( $\phi'$ ) were shown in Table 2.

This study is the first assessment of the age and growth of *A. brama*, *B. bjoerkna*, *C. gibelio*, *R. rutilus* and *V. vimba* in the lower Sakarya River. The growth parameters ( $L_{\infty}$ ,  $K$ ,  $t_0$ ,  $\phi'$ ) studied by different authors are given from other water areas (Table 3).

It was determined that the age composition of *A. brama* individuals tend to 2<sup>+</sup>-9<sup>+</sup> ages. The age composition results of different researches were given in Table 4 for *A. brama*. Asymptotic length value was higher when compared to previous studies for *A. brama* (Table 3). In contrast,  $k$  value is lower than other studies. According to the growth performance index results, it can be said that *A. brama* showed average development in conditions of the lower Sakarya River.

The maximum age of *B. bjoerkna* in this study was 10<sup>+</sup> years which is higher than that of reported studies in Table 4. These differences may be due to the variations in sampling method and period, potential aging errors, and overfishing. The  $L_{\infty}$  value obtained in this study was higher than that in earlier researches. In the present study, the  $k$  value was found to be lower than that found by other researchers (Table 3). Ma, Xie, Huo, Yang & Huang, (2010) reported that different size distributions in different study may be the causes of differences among all of the estimated parameters.

The age composition of *C. gibelio* individuals was between 1<sup>+</sup>-7<sup>+</sup> in the present study. Some differences were observed in age groups of *C. gibelio* when compared to previous researches

**Table 1.** Mean total length (ML, cm), mean weight (W, g), number of sample (n) and standard error (SE) for different age groups of five fish species.

		Age										
		1+	2+	3+	4+	5+	6+	7+	8+	9+	10+	
<i>A. brama</i>	<b>n</b>		5	14	37	55	17	8	2	3		
	<b>ML±SE</b>		18.3±2.32 (14.3-20.3)	21.2±1.30 (19.1-23.5)	24.6±1.88 (21.1-28.0)	30.0±1.17 (26.7-31.9)	33.1±0.73 (31.8-34.6)	36.1±1.75 (34.1-39.6)	39.1±0.51 (38.8-39.5)	41.4±1.16 (40.3-42.6)		
	<b>MW±SE</b>		66.92±20.21 (33.35-87.52)	104.64±23.91 (69.02-145.0)	167.60±38.44 (101.8-232.8)	299.39±46.61 (194.9-442.2)	416.40±53.25 (348.2-557.8)	560.22±73.34 (484.4-717.5)	762.81±20.49 (737.3-748.3)	872.23±45.03 (820.3-900.2)		
<i>B. bjoerkna</i>	<b>n</b>	32	43	37	82	91	80	59	71	44	8	
	<b>ML±SE</b>	11.3±1.23 (6.2-12.7)	13.9±0.83 (12.5-15.6)	16.4±0.84 (15.1-17.6)	18.3±0.76 (16.7-19.4)	19.6±0.57 (17.0-20.5)	21.1±0.60 (18.5-22.5)	22.3±0.53 (20.2-23.5)	23.9±0.82 (21.1-25.2)	26.1±0.85 (22.7-27.4)	28.3±1.14 (27.1-30.4)	
	<b>MW±SE</b>	17.82±5.15 (3.15-30.49)	30.58±6.88 (16.71-46.98)	54.39±13.68 (31.35-85.89)	80.40±10.58 (53.37-106.1)	93.86±12.54 (71.45-138.0)	115.11±18.35 (76.47-168.3)	139.45±21.65 (87.18-188.1)	177.85±31.16 (121.3-246.2)	236.59±31.70 (185.4-317.6)	302.43±46.49 (198.6-347.4)	
<i>C. gibelio</i>	<b>n</b>	45	51	15	30	23	12	3				
	<b>ML±SE</b>	12.7±0.99 (9.3-13.7)	15.9±1.64 (13.7-19.5)	21.3±0.99 (19.6-22.7)	24.3±0.90 (22.6-25.9)	26.6±0.78 (24.9-28.1)	28.8±0.91 (27.8-30.9)	31.8±0.54 (31.4-32.4)				
	<b>MW±SE</b>	40.37±8.97 (13.76-55.10)	77.10±24.92 (42.59-137.6)	191.76±37.34 (130.1-276.5)	254.18±49.50 (183.1-434.1)	325.37±61.90 (232.7-463.8)	425.02±73.72 (298.6-557.6)	499.65±80.76 (448.5-592.7)				
<i>R. rutilus</i>	<b>n</b>	25	32	28	13	15	12	13	8	6		
	<b>ML±SE</b>	12.4±0.57 (11.2-13.4)	15.3±1.35 (13.4-17.8)	18.1±1.62 (14.5-20.3)	20.6±0.78 (18.8-21.4)	22.9±1.23 (20.6-24.0)	25.1±1.35 (23.2-26.9)	27.2±1.47 (24.8-29.6)	29.1±1.26 (27.6-30.9)	30.4±1.26 (28.7-32.2)		
	<b>MW±SE</b>	22.79±3.53 (14.46-30.28)	34.34±10.94 (23.44-64.25)	62.76±20.22 (26.25-96.67)	105.89±27.48 (60.53-158.9)	128.03±20.55 (92.02-179.0)	166.25±44.87 (108.1-248.8)	218.58±55.49 (122.6-305.4)	295.58±87.46 (167.7-449.6)	382.39±73.87 (282.1-496.5)		
<i>Vimba</i>	<b>n</b>	3	68	37	20	72	46	18				
	<b>ML±SE</b>	13.0±0.59 (12.3-13.4)	14.7±1.04 (12.7-16.8)	18.8±1.31 (15.6-20.9)	21.1±0.89 (19.2-23.3)	22.6±1.02 (20.6-24.3)	24.8±0.68 (23.1-26.5)	26.8±0.89 (25.8-29.1)				
	<b>MW±SE</b>	26.20±4.66 (20.86-29.46)	31.95±8.97 (20.79-63.63)	79.95±24.35 (32.5-135.48)	116.74±14.12 (78.19-141.9)	132.86±16.81 (93.4-182.34)	156.84±22.19 (111.8-238.2)	209.07±41.89 (172.3-322.6)				

**Table 2.** Growth parameters ( $L_{\infty}$ ,  $k$ ,  $t_0$ ) and growth performance index ( $\phi'$ ) for five fish species.

Species	N	$L_{\infty}$	$k$	$t_0$	$\phi'$
<i>Abramis brama</i>	141	92.18	0.054	-0.040	2.628
<i>Blicca bjoerkna</i>	547	69.40	0.04	-0.030	2.268
<i>Carassius gibelio</i>	179	51.09	0.114	-0.024	2.474
<i>Rutilus rutilus</i>	152	48.11	0.088	-0.023	2.307
<i>Vimba vimba</i>	264	41.74	0.104	-0.035	2.260

**Table 3.** Growth parameters ( $L_{\infty}$ ,  $k$ ,  $t_0$ ) and growth performance index ( $\phi'$ ) for five fish species studied by different authors.

Species	Location	N	$L_{\infty}$	$k$	$t_0$	$\phi'$	References
<i>Abramis brama</i>	Dąbie Lake	290	44.62*** (TL)	0.175	0.23	2.542	Kompowski, 1988
	Volvi Lake	443	50.7* (FL)	0.094	-0.41	2.383	Valoukas & Economidis, 1996
	Rubikiai Lake	209	65.7*** (SL)	0.085	0.482	2.565	Žiliukienė & Žiliukas, 2011
	Sakarya River	141	92.18*** (TL)	0.054	-0.01	2.628	This study
<i>Blicca bjoerkna</i>	Berounka River		23.4*** (SL)	0.270	-0.27	2.169	Hanel, 1991
	Balaton Lake	127	35.9*** (SL)	0.098	-0.639	2.101	Specziár et al., 1997
	Sapanca Lake	350	31.91* (TL)	0.122	-1.087	2.10	Okgerman et al., 2012
	Ladik Lake	434	32.85*** (FL)	0.11	-2.64	2.074	Yilmaz et al., 2015
	Sakarya River	547	69.4*** (TL)	0.04	-0.02	2.268	This study
<i>Carassius gibelio</i>	Lysimachia Lake		32.5*** (FL)	0.282	-0.51	2.47	Leonardos et al., 2001
	Egirdir Lake	616	33.3*** (FL)	0.346	-0.302	2.58	Balik et al., 2004
	Aksu River	128	36.86*** (TL)	0.244	-0.791		Innal, 2012
	Seyhan River	317	32.30*** (TL)	0.307	-0.526	2.505	Ergüden, 2015
	Sakarya River	177	51.09*** (TL)	0.11	-0.02	2.458	This study
<i>Rutilus rutilus</i>	Volvi Lake	233	33.3*** (TL)	0.081	-1.30	1.95	Papageorgiou, 1979
	Berounka River		28.5*** (SL)	0.169	-0.17	2.14	Hanel, 1991
	Balaton Lake	112	31.9*** (SL)	0.160	0.026	2.21	Specziár et al., 1997
	Sapanca Lake	136	31.87** (TL)	0.195	-0.034	2.297	Okgerman et al., 2009
	Sakarya River	152	48.11*** (TL)	0.09	-0.02	2.318	This study
<i>Vimba vimba</i>	Berounka River		27.8*** (SL)	0.212	-0.22	2.214	Hanel, 1991
	Caspian Sea coast	845	26.1*** (FL)	0.280	-0.65	2.280	Chaichi et al., 2011
	Sapanca Lake	217	24.70* (FL)	0.205	-1.464	2.097	Okgerman et al., 2011
	Sakarya River	264	41.74*** (TL)	0.10	-0.04	2.241	This study

\*Female, \*\*Male, \*\*\*Combined

(Table 4). These differences may be due to the sampling method, fishing activity, feeding habitats, population density and the ecological conditions of water bodies.

The ages and lengths of the *R. rutilus* ranged between 1<sup>+</sup> - 9<sup>+</sup> years, 11.2 to 32.2 cm in the lower Sakarya River (Table 1). In other research on *R. rutilus*, age distribution was reported to be 1<sup>+</sup> - 6<sup>+</sup> (Sedaghat & Hoseini, 2012) in the Southern Caspian Sea, 1<sup>+</sup> - 4<sup>+</sup> in Seyhan Dam Lake (Ergüden, Ergüden, & Göksu, 2008). Due to the maximum size obtained in the sampling, asymptotic length value calculated for *R. rutilus* was found higher compared to the research in Table 3.

Despite wide distribution of *V. vimba* individuals, information on the biology of this species in Turkey is scarce. The age composition

of this species was between 1<sup>+</sup> - 7<sup>+</sup> in the present study. The growth rate for length and weight in this research was generally high in comparison with populations from other studies (Table 4). The growth performance index of *V. vimba* ( $\phi'$ ) in the lower Sakarya River was similar to that previously reported, apart from 2.097 ( $\phi'$ ) *V. Vimba* caught in the Sapanca Lake (Okgerman, Elp, & Yardımcı, 2011). The differences in the growth of *Vimba* between regions might have been because of the ecological conditions of the Sakarya River, competition for food between *Vimba* and the other fish species and differences on condition, length, age, sex, and gonadal development of *V. vimba* (Ricker, 1975).

Growth can be evaluate when age and size information are combined. Growth provides us with some indication of resource utilization and the effectiveness of our management strategies.

**Table 4.** Mean length and mean weight for different age groups of five fish species studied by different authors.

Species	Location	Sex	Age										References	
			1+	2+	3+	4+	5+	6+	7+	8+	9+	10+		
<i>Abramis brama</i>	Solina dam reservoir	TL				25.9	30.8	33.5	34.0	35.2	36.9	38.1	Epler et al., 2006	
		W				214	364	457	457	496	548	598		
	Rubikiai Lake	SL	5.6	10.0	13.6	17.0	21.5	24.3	27.6	30.2	33.2	35.9	Žiliukienė & Žiliukas, 2011	
		W	3	18	50	97	196	296	403	578	767	957		
	Ladik Lake	FL	12.72	16.14	18.08	23.9	28.75	34.09	39.53	42.16			Yilmaz et al., 2015	
		W	32.41	65.14	95.54	258.3	418.6	749.2	1167	1435.2				
	Sakarya River	TL		18.3	21.2	24.6	30.0	33.1	36.1	39.1	41.4		This study	
		W		66.92	104.64	167.60	299.39	416.40	560.22	762.81	872.23			
	<i>Billica bjoerkna</i>	Balaton Lake	SL	8.4	10.9	13.0	15.0	16.8	18.8	20.5	22.0			Specziár et al., 1997
			W	12.0	28.4	50.9	79.9	117	166	220	274			
Sapanca Lake		TL	6.9		12.99	14.96	16.73	17.80	20.01	20.77	23.2		Okgerman et al., 2012	
		W	3.35		23.32	39.23	58.28	70.02	109.73	110.0	159.4			
Uluabat Lake		FL	8.35	10.55	12.34	13.80	15.32	16.31	17.85				Şaşı & Berber, 2012	
		W	9.56	13.73	31.02	80.17	90.32	104.93	122.68					
Aras Dam Lake		TL	17.07	20.14	21.18	23.49	23.65						Jamali et al., 2015	
		W	58.7	112.2	126.1	157.9	161.7							
Sakarya River		TL	11.3	13.9	16.4	18.3	19.6	21.1	22.3	23.9	26.1	28.3	This study	
		W	17.82	30.58	54.39	80.40	93.86	115.11	139.45	177.85	236.59	302.43		
<i>Carassius gibelio</i>	Egirdir Lake	FL	11.9	18.1	22.9	25.5	27.4	29.6					Balik et al., 2004	
		W	42.0	145.2	297.0	451.4	602.1	857.5						
	Aksu River	TL	12.16	17.91	21.19	24.18	26.93	29.43					İnnal, 2012	
		W	40.71	109.41	179.43	285.38	452.21	540.3						
	Seytler Reservoir	FL	15.35	17.74	21.02	23.79	25.56				31.75		Bulut et al., 2013	
		W	46.15	130.6	214.11	300.38	348.46				755.4			
	Rozov Klade-nets Reservoir	SL	11.64	15.78	17.86	19.78	21.48	23.91					Zhelev et al., 2015	
		W	42.82	90.79	118.24	160.55	198.26	263.40						
	Sakarya River	TL	12.7	15.9	21.3	24.3	26.6	28.8	31.8				This study	
		W	40.37	77.10	191.76	254.18	325.37	425.02	499.65					

**Table 4.** Continue.

Species	Location	Sex	Age										References
			1+	2+	3+	4+	5+	6+	7+	8+	9+	10+	
<i>Rutilus rutilus</i>	Volvi Lake	TL	7.6	11.3	12.3	13.9	15.1	16.0	17.2	18.1	19.3	20.5	Papageorgiou, 1979
		W	2.93	13.59	19.30	24.86	39.98	48.74	63.85	71.92	77.24	101.35	
	Solina Reservoir	TL			15.8	19.9	21.3	22.3	23.5	25.0			Epler et al., 2005
		W			47.3	101.2	125.7	152.5	201.5	210.0			
	Seyhan Dam Lake	FL	14.72	17.40	19.25	22.04							Ergüden et al., 2008
		W	47.23	75.12	94.50	185.62							
	Sapanca Lake	TL	7.25	12.79	16.26	18.67	20.10	21.66	24.13	28.58	30.97		Okgerman et al., 2009
		W	5.19	25.38	47.56	81.14	99.01	122.96	183.59	315.34	412.53		
	Sakarya River	TL	12.4	15.3	18.1	20.6	22.9	25.1	27.2	29.1	30.4		This study
		W	22.79	34.34	62.76	105.89	128.03	166.25	218.58	295.58	382.39		
<i>Vimba vimba</i>	Sarıyar Dam Lake	FL		10.45	14.61	16.98	19.12	20.97	23.12				Ekmekci & Erk'akan, 1992
		W		13.50	42.02	74.55	112.05	161.07	235				
	Kirmir Stream	FL	10.62	13.01	15.01	17.78	19.54						Tutucu, 2002
		W	15.64	31.89	51.52	88.70	124.42						
	Barycz River	TL	6.2	12.2	17.9	23.5							Łuszczek-Trojnar et al., 2008
		W	2.25	17.1	67.6	145.9							
	Sapanca Lake	FL		12.62	14.58	17.03	17.87	19.43					Okgerman et al., 2011
		W		27.47	46.49	71.38	82.75	103.58					
	Sakarya River	TL	13.0	14.7	18.8	21.1	22.4	24.8	26.8				This study
		W	26.20	31.95	79.95	116.74	132.86	156.84	209.07				

When we evaluate age and growth in combination, the relationship between population size and biomass can be easier to understand. This understanding is the basis of modern fisheries resource allocation and management.

## CONCLUSION

In conclusion, this study provides basic information on age and growth of *A. brama*, *B. bjoerkna*, *C. gibelio*, *R. rutilus* and *V. vimba* living in the lower Sakarya River. The results of this study are useful for evaluating the relative condition of fishery managements and stock assessment in the Sakarya River. Also, this study will contribute to further scientific studies in the same area.

**Acknowledgements:** We would like to thank Dr. İrem KÖSE REİS and all fishermen for their help.

**Ethics Committee Approval:** Legal research ethics committee approval permissions for the survey were obtained from the Adnan Menderes University, Animal Experiments Local Ethics Committee.

**Financial Disclosure:** This study was funded by Muğla Sıtkı Koçman University, Scientific Research Project Office with Project number 17/073.

**Conflict of Interest:** The authors have no conflicts of interest to declare.

## REFERENCES

- Bulut, S., Ramazan, M., Algan, B., Özbek, M., Bülent, U. & Konuk, M. (2013). Several growth characteristics of an invasive Cyprinid fish (*Carassius gibelio* Bloch, 1782). *Notulae Scientia Biologicae*. [CrossRef]
- Campana, S. E. & Thorrold, S. R. (2001). Otoliths, increments, and elements: keys to a comprehensive understanding of fish populations? *Canadian Journal of Fisheries and Aquatic Sciences*. [CrossRef]
- Chaichi, A., Vosoughi, G., Kaymaram, F., Jamili, S. & Fazli, H. (2011). Population dynamics of *Vimba vimba* persain Iranian waters of the Caspian Sea. *Cybiurn*, 35(3), 237-243.
- Chugunova, N. I. (1963). *Age and Growth Studies in Fish*. Israel Prog. Sci. Transl. No: 610, Natl. Sci. Found Washington, D.C. pp 132.
- Eschmeyer, W. N., Fricke R. & Van der Laan, R. (Eds.), (2017). Catalog of Fishes. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>
- Ekmekçi, F. G. & Erk'akan, F. (1992). Some of the Growth and Reproduction Properties of *Vimba vimba tenella* (Nordmann, 1840) in Sarıyar Dam Lake. *Doga-Tr J of Zoology*, 16, 323-341.
- Emiroğlu, O. S. M. (2011). Alien fish species in upper Sakarya River and their distribution. *African Journal of Biotechnology*, 10(73), 16674-16681. [CrossRef]
- Epler, P., Popek, W., Luszczek-Trojnar, E., Drag-Kozak, E., Szczerbik, P. & Socha, M. (2005). Age and growth rate of the roach (*Rutilus rutilus* L.) from the Solina and the Tresna (Zywieckie Lake) dam reservoirs. *Acta Scientiarum Polonorum*, 4, 59-70.
- Epler, P., Lauszczek-Trojnar, E., Drag-Kozak, E., Szczerbik, P., Popek, W. & Socha, M. (2006). Age and growth of bream (*Abramis brama* L.) in the Solina, Tresna and Roznow dam reservoirs, *Acta Scientiarum Polonorum*, 5(1), 45-56.
- Ergüden, S. A., Ergüden, D. & Göksu, M. Z. L. (2008). Growth Properties of Roach (*Rutilus rutilus* L., 1758) in Seyhan Dam Lake (Adana). *Journal of FisheriesSciences.com*. [CrossRef]
- Ergüden, S. A. (2015). Age and Growth Properties of Prussian Carp, *Carassius gibelio* (Bloch, 1782) Living in the Middle Basin of Seyhan River in Adana, Turkey. *Pakistan Journal of Zoology*, 47(5), 1365-1371.
- Hanel, L. (1991). Growth of four cyprinid fishes in the river Berounka (Czech Republic). *Zivocisna Vyroba*, 36, 929-937.
- İnnal, D. (2012). Age and growth properties of *Carassius gibelio* (Cyprinidae) living in Aksu river Estuary (Antalya-Turkey). *Review of Hydrobiology*, 5(2), 97-109.
- Jamali, H., Eagderi, S., Esmailzadegan, E. & Patimar, R. (2015). Age, growth and some biological characteristics of Silver bream (*Blicca bjoerkna* L., 1758) (Cyprinidae) from Aras Dam Lake in Northwest of Iran. *International Journal of Aquatic Biology*.
- Kahraman, A. E., Göktürk, D. & Aydın, E. (2014). Length-Weight Relationships of Five Fish Species from the Sakarya River, Turkey. *Annual Research & Review in Biology*. [CrossRef]
- Kompowski, A. (1988). Growth rate of freshwater bream, *Abramis brama* (L., 1758), in Lake Dąbie and the Szczecin Lagoon. *Acta Ichthyologica et Piscatoria*. [CrossRef]
- Korkmaz, A. Ş. & Zencir Tanır, Ö. (2016). Fish Species Biodiversity in Kirmir Stream of Sakarya River, *Journal of Limnology and Freshwater Fisheries Research*. [CrossRef]
- Kottelat, M. & Freyhof, J. (2007). *Handbook of European freshwater fishes*. Publications Kottelat, Cornol and Freyhof, Berlin. 646 p.
- Lagler, K.F. (1966). *Freshwater fishery biology*. W.M.C. Brown Company Publishers Dubuque, Iowa.
- Leonardos, I., Katharios, P. & Charisis, C. (2001). Age, growth and mortality of *Carassius auratus gibelio* (Linnaeus, 1758) (Pisces: Cyprinidae) in Lake Lysimachia. p. 257-259. In Proceedings of the Tenth Ichthyological Congress, 18-22 October 2001, Chania, Greece.
- Luszczek-Trojnar, E., Drag-Kozak, E., Kleszcz, M., Popek, W. & Epler, P. (2008). Gonadal maturity in vimba (*Vimba vimba* L.) raised in carpponds. *Journal of Applied Ichthyology*. [CrossRef]
- Ma, B. S., Xie, C. X., Huo, B., Yang, X. F. & Huang, H. P. (2010). Age and growth of a long-lived fish *Schizothorax o'connori* in the Yarlung Tsangpo River, Tibet. *Zoological Studies*, 49(6), 749-759.
- Memiş, D., Tosun, D. D., Yamaner, G., Tunçelli, G., & Gessner, J. (2019). Present Status of Sturgeon in the Lower Sakarya River in Turkey. *Aquatic Research*, 2(2), 53-60. [CrossRef]
- Nelson, J. S. (1994). *Fishes of the World*. New York: John Wiley & Sons. ISBN 0471547131.
- Okgerman, H., Oral, M. & Yigit, S. (2009). Biological aspects of *Rutilus rutilus* (roach) in Sapanca Lake (Turkey). *Journal of Animal and Veterinary Advances*.
- Okgerman, H., Elp, M. & Yardimci, C.H. (2011). Growth, the length-weight relationship, and reproduction in *Vimba vimba* L. 1758) sampled from an oligo-mesotrophic lake in northwest Anatolia (Turkey). *Turkish Journal of Zoology*. [CrossRef]
- Okgerman, H., Elp, M. & Atasagun, S. (2012). The growth and reproduction of White bream (*Blicca bjoerkna* L. 1758) in an oligo-mesotrophic lake in northwest Anatolia (Sapanca, Turkey). *Turkish Journal of Biology*. [CrossRef]
- Ölmez, M. (1992). A study on the population dynamics on fishes in the Sakaryabaşı region of the upper Sakarya basin. [PhD thesis]. Ankara University. 228 s.
- Papageorgiou, N. K. (1979). The length weight relationship, age, growth and reproduction of the roach *Rutilus rutilus* (L.) in Lake Volvi. *Journal of Fish Biology*. [CrossRef]
- Pauly, D. & Munro, J. L. (1984). Once more on the comparison of growth in fish and invertebrates. *Fishbyte (Philippines)*, 1, 21-22.
- Reis, İ., Cerim, H., & Ates, C. (2019). Length-weight relationship of 13 fishspecies from the Lower Sakarya River, Turkey. *Aquatic Sciences and Engineering*. [CrossRef]
- Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish populations. *Bulletin Fisheries Research Board of Canada*, 191, 1-382.

- Şaşı, H. & Berber, S. (2012). Age, growth and some biological characteristics of White bream (*Blicca bjoerkna* L., 1758) in Uluabat Lake, in northwestern of Anatolia. *Asian Journal of Animal and Veterinary Advances*. [CrossRef]
- Sedaghat, S. & Hoseini, S. A. (2012). Age and Growth of Caspian Roach, *Rutilus rutilus caspicus* (Jakowlew, 1870) in Southern Caspian Sea, Iran. *World Journal of Fish and Marine Sciences*.
- Şengörür, B. & İsa, D. (2001). Factor analysis of water quality observations in the Sakarya River. *Turkish Journal of Engineering and Environmental Sciences*, 25(5), 415-426.
- Shiklomanov, I. A. (1999). *World water resources and their use*. St. Petersburg, Russia: State Hydrological Institute / UNESCO.
- Specziár, A., Tólg, L. & Bíró, P. (1997). Feeding strategy and growth of cyprinids in the littoral zone of Lake Balaton. *Journal of Fish Biology*. [CrossRef]
- Tutucu, S. (2002). Investigated of Some Biological Characteristics of vimba (*Vimba vimba tenella* (Nordmann, 1840)) in Kirmir Stream of Sakarya River. Natural Sciences Institute. M.Sc. thesis, Gazi University, pp 53.
- Valoukas, V.A. & Economidis, P.S. (1996). Growth, population composition and reproduction of bream *Abramis brama* (L.) in Lake Volvi, Macedonia, Greece. *Ecology of Freshwater Fish*. [CrossRef]
- Von Bertalanffy, L. (1957). Quantitative laws in metabolism and growth. *The Quarterly Review of Biology*. [CrossRef]
- Yılmaz, S., Yazıcıoğlu, O., Yazıcı, R. & Polat, N. (2015). Age, Growth and Reproductive Period of White Bream, *Blicca bjoerkna* (L., 1758) in Lake Ladik, Turkey. *Journal of Limnology and Freshwater Fisheries Research*. [CrossRef]
- Zhelev, Z., Boyadzhiev, P. & Angelov, M. (2015). Analysis of Size-Age, Sexual Structure and Condition of Populations of *Carassius gibelio* (Pisces: Cyprinidae) from two Water Basins in Galabovo Region (Southern Bulgaria). *Trakia Journal of Sciences*. [CrossRef]
- Žiliukienė, V. & Žiliukas, V. (2011). Growth rate of freshwater bream (*Abramis brama* (L.)) in Lake Rubikiai (Lithuania). *Acta Zoologica Lituana*. [CrossRef]