

## A new *Marasmius* on *Castanea sativa* from Turkey

MUSTAFA IŞILOĞLU<sup>1</sup>, HAKAN ALLI<sup>1</sup>, M. HALİL SOLAK<sup>2</sup>  
& ROY WATLING<sup>3</sup>

isiloglu48@gmail.com, hakanalli@gmail.com, halilsolak@yahoo.com  
& caledonianmyc@compuserve.com

<sup>1</sup>Muğla University, Faculty of Science and Arts, Biology Department  
48170, Muğla, Turkey

<sup>2</sup>Muğla University, Ula Ali Koçman Vocational High School, Program of Fungi  
48640 Ula, Muğla, Turkey

<sup>3</sup>Caledonian Mycological Enterprises  
Edinburgh, Scotland, United Kingdom

**Abstract** — *Marasmius castaneophilus*, a new agaric (*Marasmiaceae*, *Agaricales*, *Basidiomycota*) growing on sweet chestnut husks in Turkey, is described and illustrated. Its taxonomic position within the genus is discussed.

**Key Words** — new taxa, biodiversity, Turkish macromycota

### Introduction

*Marasmius* is a very large genus with as many as 1695 taxa (Anon 2008), even after being critically circumscribed by the transfer of some taxa to *Collybia*, *Gymnopus*, *Marasmiellus* and *Setulipes*. It is predominantly tropical in distribution (Singer 1986), with forty-two species now known from Europe (Antonin & Noordeloos 1993).

The species within this restricted concept are often highly specific as to their host requirements. This specificity is not only to the host, but to a particular part of the host colonized, whether it be leaf lamina, midrib or petiole (Watling 1982). Traditionally the genus was split into those with pileipellis in the form of Rameales-structure, viz. cutis of diverticulate hyphae, and setiform stipe (sect. *Androsacei* Kühner, = *Setulipes* Antonín) and those with a hymeniform pileipellis (other *Marasmius* sections). As traditionally accepted, the genus

\* corresponding author

This article is dedicated to Dr. Fadime Yılmaz (1971-2005), Muğla University, who made many contributions to our knowledge of Turkish macrofungi (including members of the genus *Marasmius*) but, sadly, died in a car accident in 2005.

*Marasmius* contains a group of white or pale coloured, small fruitbodies usually characterized as having an insititious stipe and well-developed cystidia and referred to sect. *Epiphylli* Kühner. European representatives of this section include *M. epiphylloides*, *M. epiphyllus*, *M. hellebori-corsici*, *M. saccharinus*, *M. setosus*, and *M. tremulae* (= *M. favrei*). However, molecular studies place this section not in *Marasmius* s. str., but rather close to the *Physalacriaceae* (Owings & Desjardin 1997, Moncalvo et al. 2002, Wilson & Desjardin 2005), restricting the genus *Marasmius* only to representatives of the sections *Marasmius*, *Sicci*, *Globulares*, *Hygrometrici*, *Neosessiles* and *Leveilleani*. Recent studies in Turkey have encountered an additional member of the group related to *M. epiphylloides* that is apparently confined to colonizing the husks of *Castanea sativa*. The species described formally below raises the total number of members of *Marasmius* in Turkey to 23 (Solak et al. 2007).

### Taxonomic description

*Marasmius castaneophilus* Işıloğlu, Allı, Solak & Watling, sp. nov.

FIGS. 1–5

MYCOBANK MB511864

*Pileus* 4–4.5 mm convexus vel plano-convexus prostromo plano, interdum depressus, albus, radiato-striatus vel radiato-sulcatus ad marginem interdum undulato-lobatus. Lamellae lato-adnatae, distantes, furcatae, plicatae, albae. Stipes 20–30 × 0.5–1 mm., filiformes vulgo flexuosus, glabrus, ad apicem hyalinum prostromo ad basim rufo-brunneum, vulgo senecutate infra atro-ferrugineus. Caro albo. Odor, sapor nullus. Sporae 10–12 × 4–5 µm, ellipsoideo-oblongae vel fusiformae, laevae, hyalinae, inamyloideae. Basidia 45–50 × 4–6 µm, 4-sporigera. Cystidia faciei lamellarum fusiformia vel clavata, tenuio-tunicata; cystidia aciei lamellarum 1) 30–40 × 5–8 µm, subfusiformia vel lageniformia et 2) 15–35 × 15–20 µm, clavata vel pyriformia ornamentatae. Cellulae cuticulae pilei hymeniformae 1) 15–18 × 8–10 µm, clavata, tenuio-tunicata – ‘rotalis-typus’ et 2) 20–27 × 4–5 µm, fusiformiae vel lageniformiae, tenuio-tunicata, laevae. Hyphae fibulatae. Trama inamyloideae. Ad *Castanea sativa* cupulus. Turkey; İzmir, Ödemiş Typus Ha 2842 in E.

PILEUS 4–4.5 mm, convex to plano-convex then irregularly applanate, sometimes depressed, white, radially striate or grooved with undulating margin (FIG. 1). GILLS broadly adnate, poorly developed, distant, vein-like, forked, plicate and without lamellulae, always white. STIPE 20–30 × 0.5–1 mm., hair-like, central, rarely eccentric, glabrous, smooth, twisted when dry, stiff and tough, hyaline when young, reddish brown below when old, blackish brown when dry. FLESH very thin and white. SMEL AND TASTE indistinct. SPORE PRINT White. SPORES 10–12 × 4–5 µm, fusiform to narrowly elliptic with a long apiculus (FIG. 2), smooth, thin-walled, colourless in alkali, inamyloid. BASIDIA 45–50 × 4–6 µm, clavate, hyaline, thin-walled, 4-spored, with clamps. BASIDIOLES 35–40 × 4–5 µm, clavate, with clamps. PLEUROCYSTIDIA 40–70 × 9–12 µm, fusiform to clavate, thin-walled and hyaline (FIG. 3). CHEILOCYSTIDIA of two types; 1. type: 30–40 × 5–8 µm subfusiform to lageniform with long neck, thin-walled (FIG.



FIG. 1. *Marasmius castaneophilus* (from holotype; photo by H.Alli)

4). 2. type:  $15\text{--}35 \times 15\text{--}20 \mu\text{m}$ , broom cells of the rotalis-type, broadly clavate to pyriform with projections. PILEIPELLIS hymeniform, a mixture of three types of elements. 1)  $15\text{--}18 \times 8\text{--}10 \mu\text{m}$ , clavate, thin-walled, rotalis-type cells with long projections 2) smooth clavate cells and 3)  $20\text{--}27 \times 4\text{--}5 \mu\text{m}$ , fusiform or lageniform, thin-walled and smooth (FIG. 5); the former the most common. CLAMP-CONNECTIONS present.


CHEMICAL REACTIONS. No part of basidiome dextrinoid or amyloid.

HABITAT. In clusters on dead husks of *Castanea sativa*.

SPECIMEN EXAMINED. Turkey; İzmir, Ödemiş, Elmabağı village, in natural forest of mixed *Pinus nigra* and *Castanea sativa*, 11 November 2007, Ha 2842. Holotype: in E.

## Discussion

*Marasmius castaneophilus* is easily recognized in the field by its unique habitat on sweet chestnut husk, pure white pileus surmounting a stipe that becomes red brown upwards with age, lanceolate pileocystidia, and large basidiospores. Following Antonín & Noordeloos (1993), this species belongs to sect. *Epiphylli* subsect. *Epiphyллоidei* Singer, whose taxa are characterized by having pilepellis composed of a mixture of smooth cells and rotalis-type broom cells, absence of distinct collarium, marasmoid or collybioid basidiomes, filiform and insititious



FIGS. 2–5. *Marasmius castaneophilus* (from holotype).  
2. spores, 3. pleurocystidia, 4. cheilocystidia, 5. pileipellis

stem, white or pale pileus, vein-like and reduced gills, and non-dextrinoid hyphae. It is closely related to *M. epiphylloides* (Rea) Sacc. & Trotter and can be distinguished confidently both morphologically and microscopically. *Marasmius castaneophilus* has a longer hair like stem, larger basidiospores, and two types of cheilocystidia. It also differs by its association with sweet chestnut husks.

#### Acknowledgements

The authors would like to express their thanks to Drs. Vladimír Antonín (Moravian Museum, Brno, Czech Republic) and Machiel E. Noordeloos (Rijksherbarium, Leiden) for reviewing the manuscript, to Ferah Yılmaz (Muğla University) for helping to collect the specimen and to Rabia Melike Işıloğlu (Muğla University) for helping to draw figures. This study was supported financially by Tübitak, The Scientific and Technical Research Council of Turkey (project no: TBAG-102T106).

### Literature cited

- Anon 2008. <http://www.mycobank.org/Mycotaxo.aspx>.
- Antonin V, Noordeloos ME. 1993. A Monograph of *Marasmius*, *Collybia* and Related Genera in Europe. Part 1: *Marasmius*, *Setulipes*, and *Marasmiellus*. Libri Botanici 8. IHW-Verlag, Eching.
- Moncalvo JM, Vilgalys R, Redhead SA, Johnson JE, James TY, Aime MC, Hofstetter V, Verduin SJV, Larsson E, Baroni TJ, Thorn RG, Jacobsson S, Clémenton H, Miller OK. 2002. One hundred and seventeen clades of euagarics. *Molecular Phylogenetic and Evolution* 23: 357–400.
- Owings P, Desjardin DE. 1997. A molecular phylogeny of *Marasmius* and selected segregate genera. *Inoculum* 48(3): 29.
- Singer R. 1986. *Agaricales* in *Modern Taxonomy*. 4th ed. J. Cramer, Vaduz.
- Solak MH, Işıoğlu M, Kalmış E, Allı H. 2007. Macrofungi of Turkey, Checklist. Üniversiteler ofset, İzmir.
- Watling R. 1982. Taxonomic Status and Ecological Identity in the *Basidiomycetes*. In *Decomposer Basidiomycetes: their Biology and Ecology* (eds. Frankland, Hedger, Swift). Pitman Press, Bath, UK.
- Wilson AW, Desjardin DE. 2005. Phylogenetic relationships in the gymnopoid and marasmioid fungi (*Basidiomycetes*, euagaric clade). *Mycologia* 97(3): 667–679.

