

Hepatoprotective Effects of *Hertia cheirifolia* Butanolic Extract and Selenium against CCl₄-induced Toxicity in Rats

Abstract

Background: *Hertia cheirifolia*, a traditional plant endemic to both Tunisia and Algeria, is used for the treatment of various disorders. This study investigates the antioxidant and protective effects of *H. cheirifolia* butanolic extract (BEHC) alone and combined with selenium (Se) against carbon tetrachloride (CCl₄)-induced liver damage in rats. **Experimental Procedure:** Thirty male Wistar rats were randomly divided into six groups: (1) normal control, (2) hepatotoxic control, (3) positive control received silymarin 100 mg/kg body weight (bw), (4) BEHC (100 mg/kg bw), (5) BEHC (400 mg/kg bw), and (6) BEHC (400 mg/kg bw) + Se (0.3 mg/kg bw) once daily for 14 consecutive days, followed by hepatotoxicity induction with CCl₄ in olive oil 0.6 mL/kg bw intraperitoneally. Some biochemical and oxidative stress parameters were investigated. Quantity and quality of phenolics in BEHC were determined by spectrophotometer and high-performance liquid chromatography with diode-array detection (HPLC-DAD) analysis, respectively. **Results and Conclusion:** BEHC contained high amounts of total phenolics and flavonoids where seven compounds were identified. The pretreatment with BEHC or with BEHC and Se significantly reduced the levels of plasma aminotransferases (alanine aminotransferase [AST] and aspartate aminotransferase [ALT]), alkaline phosphatase, malondialdehyde (MDA), and increasing glutathione (GSH), catalase (CAT), and superoxide dismutase (SOD) levels in hepatic tissues. In conclusion, BEHC has a potent natural antioxidant activity that can be used with Se to reduce hepatotoxicity.

Keywords: Carbon tetrachloride, hepatotoxicity in rat, *Hertia cheirifolia*, oxidative stress, selenium

Introduction

Carbon tetrachloride (CCl₄) is a haloalkane hepatotoxin generally employed as a solvent, cleaner, and degreaser both for industrial and home use.^[1] This agent proves extremely useful as an experimental model for the study of certain hepatotoxic effects and serves to estimate the efficiency of hepatoprotectants.^[2] CCl₄ is metabolized and transformed to trichloromethyl and trichloromethyl peroxy radicals through cytochrome P₄₅₀ complex; these free radicals initiate the chain reaction of lipid peroxidation, which destroys polyunsaturated fatty acids inducing damages in different organs as well as liver, kidney, testis lung, and brain.^[3]

Medicinal plants are known as an important source of new drugs. These natural resources appear interesting to develop alternative treatments. Moreover, plants are rich in antioxidants and commonly used against oxidative stress-related disorders and tissue injuries.^[4] Some traditional plants have a

substantial hepatoprotective effect against various experimental animal models.^[5] One of these important traditional hepatoprotective drugs is Silymarin (*Silybum marianum*); this flavonolignan displays important hepatoprotective effects as it prevents the penetration of hepatotoxic substances, such as CCl₄ by altering cytoplasmic membrane architecture.^[6] Selenium (Se) is an essential dietary trace component, which plays an antioxidant role because it is an integral part of various proteins with catalytic and structural functions. The nutritional deficiency of Se in humans leads to chronic degenerative disorders that could be prevented by the supplementation of Se when used alone or in combination.^[7]

In this study, we chose the species *Hertia cheirifolia* to estimate its hepatoprotective effects against CCl₄-induced injury in the liver of rat. The genus *Hertia* which belongs to the *Asteraceae* family is distributed with its 12 species in the South and North Africa and South-West Asia.^[8] In Algeria, it was found only the species *H. cheirifolia* (L).^[9,10]

How to cite this article: Menakh M, Boutellaa S, Mahdi D, Zellagui A, Lahouel M, Öztürk M. Hepatoprotective effects of *Hertia cheirifolia* butanolic extract and selenium against CCl₄-induced toxicity in rats. *J Rep Pharm Sci* 2021;10:216-24.

Mouna Menakh¹,
Saber Boutellaa^{2,3},
Djahida Mahdi^{1,4},
Amar Zellagui²,
Mesbah Lahouel⁵,
Mehmet Öztürk⁶

¹Department of Nature and Life Sciences, Faculty of Exact Sciences and Nature and Life Sciences, Oum El Bouaghi University, Oum El Bouaghi, Algeria, ²Laboratory of Biomolecules and Plant Breeding, Department of Nature and Life Sciences, Faculty of Exact Sciences, Oum El Bouaghi University, Oum El Bouaghi, Algeria, ³Department of Nature and Life Sciences, University Center Abdehfid Boussouf, Mila, Algeria, ⁴Laboratory of Animal Eco-Physiology, Department of Biology, Faculty of Sciences, Badji Mokhtar University, Annaba, Algeria, ⁵Laboratory of Molecular Toxicology, Faculty of Sciences, Jijel University, Jijel, Algeria, ⁶Department of Chemistry, Faculty of Sciences, Muğla Sıtkı Koçman University, Muğla, Turkey

Received: 05 Apr 2020

Accepted: 16 Oct 2021

Published: 17 Dec 2021

Address for correspondence:

Dr. Mouna Menakh,
Department of Nature and Life Sciences, Faculty of Exact Sciences and Nature and Life Sciences, Oum El Bouaghi University, Oum El Bouaghi 04000, Algeria.
E-mail: bout.mouna@gmail.com

Access this article online

Website:

www.jrpsjournal.com

DOI: 10.4103/jrtps.JRTPS_25_20

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

This plant, which is also known as *Othonnopsis cheirifolia*, is endemic to both Tunisia and Algeria; it is traditionally used to treat inflammatory disorders, pain of stomach, diarrhea, and to reduce hyperglycemia.^[9]

Previous studies showed that *H. cheirifolia* has considerable chemicals and biological activities.^[10] Moreover, the methanolic, ethyl acetate, and chloroformic extracts of *H. cheirifolia* were tested for their spasmolytic and anti-inflammatory activities.^[11] The antioxidant and the protective activities of *H. cheirifolia* methanol and aqueous extracts against biomolecule oxidative damages were also evaluated.^[9] Lots of studies were reported about the biological properties of essential oils from *H. cheirifolia* such as the acaricidal effects,^[12] antioxidant activity, and inhibitory properties against α -glucosidase.^[13]

This study aimed to evaluate the potential protective effects of *H. cheirifolia* butanolic extract (BEHC) alone and combined with Se against CCl₄-induced liver oxidative stress in rats. In addition, the plant extract was analyzed by high-performance liquid chromatography with diode-array detection (HPLC-DAD) and examined for its antioxidant potential.

Materials and Methods

Extract preparation

The aerial parts of *Hertia cheirifolia* (HC) were collected from Oum El Bouaghi (East of Algeria) during the flowering period (April 2015). The plant was identified by Professor Zellagui Amar and a voucher specimen was deposited in the Laboratory of Biomolecules and Plant Breeding, University of Larbi Ben Mhidi Oum Elbouaghi (Algeria) under number ZA 122. Samples were cleaned, dried in shade, and powdered using an electric milling, and then 2 kg of powdered plant was macerated three times in an ethanol or water mixture (70:30 v/v) from 48 to 72 h by renewing the solvent each time. The obtained solution was filtered, concentrated, and kept standing overnight for decantation. Using a 1-L separating funnel, we proceeded to successive liquid-liquid extractions, by four organic solvents of increasing polarity: petroleum ether, chloroform, ethyl acetate, and *n*-butanol.^[14]

Total phenolics and flavonoids contents

To determine total phenolic content in BEHC in each tube, a volume of 0.5 mL (in triplicate) for each extract was added to 2.5 mL of 10% FC reagent. After 10 min of incubation at ambient temperature, the reaction medium was alkalized with 2 mL of sodium carbonate (7.5%). All tubes were shaken and incubated for 1 h in the dark before measuring the absorbance at 760 nm by a ultraviolet (UV) spectrophotometer. The results were expressed as milligram of gallic acid equivalent per gram of extract (mg GAE/g extract).^[15]

The BEHC was also analyzed spectrometrically to determine flavonoid content using quercetin (5–20 μ g/mL) as standard. 1 mL of extract (1 mg/mL) was added to 1 mL of AlCl₃ (2%), incubated for 10 min at room temperature. Then, the absorbance was

measured at 430 nm, and the results were expressed as milligram quercetin equivalent per gram of extract (QE/g extract).^[16]

High-performance liquid chromatography analyses

The analysis of phenolic constituents present in the BEHC was performed using a Shimadzu reverse phase HPLC-DAD (Shimadzu Cooperation, Japan). The column temperature was set at 35°C. The separation was carried out on a Inertsil ODS-3 (4 μ m, 4.0 mm \times 150 mm) column and Inertsil ODS-3 guard column; mobile phases were aqueous acetic acid 0.1% (A) and methanol (B). The injected volume was 20 μ L. The separation was carried out using a diode-array detector (DAD) at 254-nm wavelength. All the samples and standards were filtered with an Agilent 0.45 μ m filter.^[17]

Free radical scavenging activity

The antiradical activity of the BEHC was measured using the purple-colored solution of 1,1-diphenyl-2-picrylhydrazyl radical (DPPH). Briefly, 39.4 mg of DPPH was dissolved in methanol to prepare 0.1 mM of solution and 4 mL of this solution was added to 1 mL of BEHC in methanol at different concentrations. The resulting solution was shaken vigorously, and after 30 min of incubation in the dark at room temperature, its absorbance was measured at 517 nm. Butylated hydroxytoluene (BHT) and butylatedhydroxyanisole (BHA) were used as standards antioxidants. The ability to scavenge the DPPH was calculated using the following formula:

$$\text{Scavenging activity \%} = [\text{Abs (control)} - \text{Abs (sample)}] / \text{Abs (control)} \times 100.$$

The results were expressed as half-maximal inhibitory concentration (IC₅₀) (μ g/mL), which represented the concentration of extract required to cause a 50% DPPH inhibition.^[18]

Experimental animals

Thirty male Wistar rats, weighing 130 \pm 13 g, were used in this study. They were purchased from the breeding division of animals at Pasteur Institute located in Algiers (Algeria), and housed in plastic cages (5 animals/cage). The animals were maintained under standard laboratory conditions of constant temperature (24 \pm 2°C), relative humidity (60%), 12 h light: 12 h dark cycle, and allowed free access water and standard pellet rat diet provided by National Livestock Food Board (Bejaia, Algeria). All experiments were performed according to the international guidelines. The study protocol was designed and approved by the Consultative Ethics Committee of the Biotechnology Research Centre in Constantine, Algeria (CCE-8-10-2014).

Experimental design

The 30 experimental rats were divided randomly into six groups, five rats in each group, and treated for 14 days^[19] as follows:

- Group 01: Normal control—rats of this group received daily normal saline solution for 14 days, and then administered

0.6 mL/kg bw of olive oil, which served as vehicle intraperitoneally on the last day of the treatment.

- Group 02: Hepatotoxic control—rats of this group received daily normal saline solution for 14 days, followed by 0.6 mL/kg bw of CCl₄ (dissolved in olive oil v/v) by intraperitoneal injection before 24 h of the sacrifice, which is clearly documented to induce hepatotoxicity in rats.^[20]
- Group 03: Standard (positive control) group—rats of this group received daily a single dose (100 mg/kg bw) of silymarin for 14 days, followed by 0.6 mL/kg bw of CCl₄ by intraperitoneal injection before 24 h of the sacrifice.
- Group 04: Based on previous studies which showed that polar extract from *H. cheirifolia* was safe and has no toxicity effects even at 2000 mg/kg.^[21] Rats of this group received daily 100 mg/kg bw BEHC for 14 days, followed by 0.6 mL/kg bw of CCl₄ by intraperitoneal injection before 24 h of the sacrifice.
- Group 05: Rats of this group received daily 400 mg/kg bw BEHC for 14 day, followed by 0.6 mL/kg bw of CCl₄ by intraperitoneal injection before 24 h of the sacrifice
- Group 06: Rats of this group received daily 400 mg/kg bw BEHC associated with 0.3 mg/kg bw of Se (as Na₂SeO₃) for 14 days followed by 0.6 mL/kg bw of CCl₄ by intraperitoneal injection before 24 h of the sacrifice.

Sample collection

At the end of the experiment, rats were anesthetized with chloroform, blood samples were collected immediately from the heart into heparinized tubes, and the livers were removed rapidly, dissected, and washed to remove excess blood and cut into pieces.

Measurement of serum biochemical markers

Blood samples were centrifuged at 2000 rpm for 10 min, and then the plasma was removed immediately and stored at -20°C. The levels of plasma total protein (TP), total bilirubin (TB), cholesterol (CHOL), triglycerides (TG), alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase (ALP) activities were measured using commercially standard kits (Spinreact, Espagne).

Preparation of liver cytosolic fraction

Approximately 1 g of liver was homogenized in 3 volume of buffer solution of phosphate-buffered saline (0.1M, pH 7.4) and KCl 1, 17%. Homogenates were centrifuged at 2000 rpm for 15 min at 4°C, and the obtained supernatant was centrifuged at 9600 rpm for 45 min at 4°C, and the resultant cytosolic fraction was used for the determination of catalase (CAT), superoxide dismutase (SOD), glutathione (GSH), and malondialdehyde (MDA) levels.^[22]

Measurements of oxidative stress indicators

CAT activity was estimated by the UV colorimetric method using H₂O₂ as substrate and enzymatic activity of each sample was measured in international units (IU)/mg of proteins.^[23] SOD enzyme activity measurement was conducted using

the oxidizing reaction of nitroblue tetrazolium (NBT). The absorbance was determined at 560 nm, and the specific activity of each sample was estimated in U/min/mg of protein.^[24]

Measurement of liver GSH was performed using a colorimetric technique based on the change of a yellow color when DTNB [5,5 dithiobis-(2-nitrobenzoic acid)] is added to compounds containing sulfhydryl groups. The absorbance was recorded at 412 nm and the results were expressed as nmol GSH/mg protein.^[25]

MDA content in liver was estimated using tetramethoxypropane as a standard. The absorbance was recorded at 535 nm and MDA levels were expressed as nmol MDA/mg protein.^[26]

Statistical analysis and calculation

All data are expressed as mean ± standard error of the mean (SEM). Testing for statistical significance was assessed by a one-way analysis of variance (ANOVA), followed by Tukey's *post hoc* test using Statistica software (version 5.1, StatSoft, France). Values of *P* < 0.05 were considered significant. Percentages of change and improvement were calculated according to the following equations:^[27]

% Change = [(mean of control – mean of treated)/mean of control] × 100%, Improvement = [(mean of disease (CCl₄) – mean of treated)/mean of control] × 100.

Results

Total phenolics and flavonoid contents

Total phenolic and flavonoid contents of *n*-butanolic extract of *H. cheirifolia* are presented in Table 1. The results showed that BEHC had a high amount of total phenolics (203.52 ± 1.81 mg GAE/g) and flavonoids (104.86 ± 0.57 mg QE/g).

High-performance liquid chromatography profiling

The results of HPLC analysis are given in Table 2. Seven compounds were detected in *n*-butanolic extract of *H. cheirifolia*. Rutin was the major compound (36.32 µg/g), followed by trans-2-hydroxycinnamic acid (5.41 µg/g), ferulic acid (3.04 µg/g), 6, 7 dihydroxycoumarin (2.28 µg/g), chlorogenic acid (2.21 µg/g), 4-hydroxybenzoic acid (0.49 µg/g), and a trace of 2, 4,-dihydroxybenzoic acid.

Free radical scavenging activity

The BEHC was found to have a higher scavenging activity of DPPH [Figure 1] with an IC₅₀ of 91.60 µg/mL as compared with that of both used standard antioxidants BHT (IC₅₀ = 22.32 µg/mL) and BHA (IC₅₀ = 5.73 µg/mL).

Effects of *Hertia cheirifolia* butanolic extract on plasma biochemical parameters and liver enzymes

It was found that TB, ALP, AST, ALT, TG, and CHOL values increased significantly in the hepatotoxic group as compared to control ones (*P* < 0.01 for TB, and *P* < 0.001 for ALP, AST, ALT, TG, and CHOL) [Figure 2] with percentages increase of 183.33%, 449.36%, 182.77%, 283.79%, 112%, and 46.66%,

Table 1: Total phenolics and flavonoids contents in *n*-butanolic extract of *Hertia cheirifolia*

Plant extract	Total phenolics (mg GAE/g)	Total flavonoid (mg QE/g)
<i>n</i> -butanolic extract	203.52 ± 1.81	104.86 ± 0.57

mg GAE/g = mg of gallic acid equivalent per g of dry plant extract, mg QE/g = mg of quercetin equivalent per g of dry plant extract, SEM = standard error of the mean

Values are mean ± SEM of triplicate determinations

Table 2: Total phenolics and flavonoids compounds quantification and qualification of *Hertia cheirifolia* butanolic extract (BEHC)

Compounds (µg/g)	Retention time (min)	BEHC
4-hydroxybenzoic acid	8.01	0.49
6,7 dihydroxycoumarin	11.62	2.28
2,4,-dihydroxybenzoic acid	15.54	tr
Chlorogenic acid	17.05	2.21
Ferulic acid	19.76	3.04
trans-2-hydroxycinnamic acid	21.98	5.41
Rutin	22.01	36.32

BEHC = *Hertia cheirifolia* butanolic extract, (tr) = trace, SEM = standard error of the mean

Values are mean ± SEM of triplicate determinations




Figure 1: Free radical scavenging activity of *n*-butanolic extract of *Hertia cheirifolia* (BEHC), BHA, and BHT. Values are mean ± SEM of triplicate determinations

respectively. However, a highly significant decrease in liver function biomarkers (ALP, AST, ALT, and TB) was observed in groups treated with silymarin and BEHC with and without Se when compared with hepatotoxic control group ($P < 0.001$ for ALP, AST, and ALT; $P < 0.01$ for TB) showing a remarkable improvement. In addition, we noticed also that the pretreatment with BEHC (400 mg/kg) +Se showed a higher ameliorative effect for ALP, AST, ALT, and TB (421%, 165.96%, 217.57%, and 166.67, respectively) which was better than that shown by silymarin (288.55%, 36.22%, 127.45%, and 147.22%, respectively) [Figure 3].

In addition, TG and CHOL values decreased significantly in groups treated with silymarin and BEHC with and without Se compared to hepatotoxic group ($P < 0.01$) [Figure 2]. Moreover, marked improvement noticed in the lipid parameters (TG and CHOL) in groups treated with BEHC (100 and 400 mg/kg) with percentage of improvement recorded 30% and 88%, respectively, for CHOL and 33 and 88% for TG, respectively. However amelioration was recorded for CHOL

and TG in BEHC (400 mg/kg) +Se-treated group (33.33% and 92%, respectively, compared to silymarin treated group ones (35% and 96%) [Figure 3].

Moreover, TP values decreased significantly in the hepatotoxic control group as compared to control ones ($P < 0.01$) with percentages of changes amounted 26%. However, the pretreatment with BEHC 100 mg/kg, BEHC (400 mg/kg), BEHC (400 mg/kg)+ Se and silymarin increased this parameter significantly when compared to hepatotoxic group and recorded percentages of amelioration 18.36%, 16.47%, 24.56%, and 17.5%, respectively [Figure 3].

Effects of *Hertia cheirifolia* butanolic extract on oxidative stress markers

Our results showed in hepatotoxic control group a significant increase in MDA levels compared with control ones ($P < 0.001$) with percentages of changes amounted to 347.87%. In contrast, pretreatment with BEHC (400 mg/kg) and silymarin showed a significant decrease in MDA levels ($P < 0.05$ for BEHC (400 mg/kg); $P < 0.01$ for silymarin. MDA production was reduced significantly ($P < 0.01$) when Se was combined with the plant extract as silymarin did. However, pretreatment with BEHC (100 mg/kg) showed an insignificant decrease in MDA levels [Figure 4]. Moreover, a highly marked improvement was observed in MDA level with improvement percentages of 248.94%, 178.72%, 227.66%, and 201.06% for groups treated with silymarin, BEHC (100 mg/kg), and BEHC (400 mg/kg) with and without Se, respectively [Figure 5].

Results in Figure 4 showed a significant decrease in the levels of CAT, GSH, and SOD in hepatotoxic group as compared with those of control ones ($P < 0.001$), and showed percentage of changes 69.9%, 72.25%, and 49.62%, respectively. These marked changes were accompanied with a high significant amelioration by increasing in the levels of CAT and GSH




Figure 2: Effects of BEHC on plasma biochemical parameters and liver enzymes. N control = normal control, CCl₄ = intoxicated control, sily+CCl₄ = standard group (silymarin 100 mg/kg bw) +CCl₄; EXT100 mg+CCl₄; EXT400 mg+CCl₄ and EXT400 mg+Se+CCl₄: groups treated with *n*-butanolic extract (100 mg/kg bw+CCl₄; 400 mg/kg bw+CCl₄; 400 mg/kg bw+selenium 0.3 mg/kg bw+CCl₄), respectively. Values are mean ± SEM, n = 5 animals in each group. *P < 0.05, **P < 0.01, and ***P < 0.001 as compared to normal control group. #P < 0.05, ##P < 0.01, and ###P < 0.001 as compared to hepatotoxic control group




Figure 3: Percentages of improvement in different biomarkers of CCl₄-intoxicated rats treated with silymarin, BEHC 100mg/kg, 400mg/kg, and BEHC 400mg/kg+selenium

in groups treated with BEHC and silymarin compared to hepatotoxic group ($P < 0.05$ for BEHC (100 mg/kg); $P < 0.001$ for silymarin, BEHC (400 mg/kg) with and without Se). SOD level was significantly elevated in silymarin, BEHC (400 mg/kg) alone and BEHC (400 mg/kg) with Se groups ($P < 0.05$) as shown in Figure 4.

We observed also that percentages improvement in CAT, GSH, and SOD of BEHC 400mg/kg+ Se (40.28%, 32.72%, and 28.84% for CAT, GSH, and SOD, respectively) were higher

than those recorded in silymarin pretreatment group (37.62%, 25.87%, and 28.53%, respectively) [Figure 5].

Discussion

Several researches were carried out about chemical composition and biological properties of *Asteraceae*, but few studies on biological activities of *H. cheirifolia* were focused. This study investigated the protective effect of BEHC and Se against oxidative damages induced by CCl₄ in rats.

The free radical scavenging activity of BEHC was tested by measurement of the capacity of this extract to scavenge the stable free radical formed in solution, by donating of a hydrogen atom or an electron.^[28] As indicated from the results, BEHC showed concentration-dependent free radical scavenging activity. Comparing our results with those in previous studies, BEHC showed higher scavenging activity of DPPH than the butanolic extract of Tunisian *H. cheirifolia* flowers and roots (210 ± 0.01 µg/mL) and (98 ± 0.006 µg/mL), respectively.^[29,30] This capacity is probably highly related to the higher polyphenols and flavonoids contents in BEHC as shown in the results. In our previous study, a variety of *in vitro* antioxidant assays were performed to determine the antioxidant status of BEHC and the results indicated that the BEHC showed an interesting antioxidant activity when compared with those of standard antioxidants in all tests, and the highest activity was recorded in the β-carotene-linoleic acid assay (IC₅₀ = 9.99 ± 0.53 µg/mL).^[31]

In fact, the HPLC analysis of BEHC showed the presence of some important flavonoids such as Rutin (quercetin-3-rhamnosyl glucoside) as the major compound, which had been reported to possess a high antioxidant and anti-inflammatory activity.^[32] In another study, Rutin represented also the major constituent in methanol extract of *H. cheirifolia*.^[21]

Liver is the main organ of detoxification, which plays a major role in diverse metabolisms and transformation xenobiotics into compounds with low toxicity and excretes them from the body. It has been reported that various toxic chemicals such as antibiotics, chemotherapeutic agents, and CCl₄ damage liver cells.^[33]

Our results showed that the administration of CCl₄ caused liver injuries in rats, as shown by the significant increase in plasma levels of AST, ALT, and PAL. It is well known that the activity of AST, ALT, and PAL enzymes in plasma reflect the dysfunctional activity of the liver, and they are mainly




Figure 4: Effects of BEHC on plasma biochemical parameters and liver enzymes. N control: normal control; CCl₄: intoxicated control; sily+CCl₄: standard group (silymarin 100 mg/kg bw) +CCl₄; EXT100 mg+CCl₄; EXT400 mg+CCl₄ and EXT400 mg+Se+ CCl₄: groups treated with *n*-butanolic extract (100 mg/kg bw+CCl₄; 400 mg/kg bw+CCl₄; 400 mg/kg bw+selenium 0.3 mg/kg bw+CCl₄), respectively. Values are mean ± SEM, n = 5 animals in each group. *P < 0.05, **P < 0.01, and ***P < 0.001 as compared to normal control group. #P < 0.05, ##P < 0.01, and ###P < 0.001 as compared to hepatotoxic control group




Figure 5: Percentages of improvement in MDA, GSH, CAT, and SOD of CCl₄-intoxicated rats treated with silymarin, BEHC 100 mg/kg, 400 mg/kg, and BEHC 400 mg/kg + selenium

used in the assessment of liver damage and dysfunction.^[34] This increase in the enzymatic activity was confirmed also by previous reports on CCl₄-induced liver damage.^[2,5,6] This result might be due to the change of the plasma membrane permeability and consequently the leakage of enzymes from the tissue to the plasma or due to the onset of liver necrosis.^[35] According to previous investigation, extracts of *H. cheirifolia* were regarded as being safe or practically nontoxic by oral route at doses greater than 400 mg/kg b w.^[36] In our study, treatment with BEHC at the dose of 100 mg/kg, and 400 mg/kg with and without Se improved the deleterious effects of CCl₄ in decreasing significantly the levels of enzymes compared with hepatotoxic control group especially. However, the pretreatment with BEHC (400 mg/kg) accompanied with Se showed higher ameliorative effect which was better than that showed by silymarin.

Se is a trace element with a great importance for health and its deficiency can lead to several diseases. It is essential for normal metabolic processes, as well as the metabolism of thyroid hormones, antioxidant defense and immune function.^[37] The protective effects of Se appeared to be mainly associated in selenoenzymes, which are known to protect several cellular components against oxidative damage.^[38] In agreement with these results; a previous study showed that Se has protective effects against hepatotoxicity induced by CCl₄.^[39]

Liver has a fundamental role in the metabolism of lipids, glucides, and proteins. This study has also revealed that the CCl₄ injection-induced liver metabolic disorders including the destruction of lipid synthesis. In fact, data presented a highly significant increase in plasma levels of CHOL and TG. The increased esterification of fatty acids, inhibition of fatty acid β -oxidation, and decreased excretion of cellular lipids could explain the rise in CHOL levels.^[40] CCl₄ stimulates the transport of acetate to liver cells that enhance CHOL synthesis. It also increases fatty acids and TG synthesis

from acetate and increases lipid esterification.^[11] In addition, inhibition lysosomal lipase activity and very low density lipoprotein (VLDL) secretion may cause accumulation of TG in the liver.^[41] However, pretreatment with BEHC and Se decrease significantly plasma levels of CHOL and TG lessening the CCl₄-induced liver injury owing to their capacity to inhibit the damaging effects of reactive oxygen species (ROS) and avoid low density lipoprotein (LDL) and cell membrane oxidation. On the contrary, this action causes the decrease of acetate transfer to liver cells and diminishing the synthesis of CHOL, free fatty acids, and TG.^[19]

Bilirubin is a metabolic waste product formed from the destruction of aged or abnormal erythrocytes.^[42] Significant elevated level of plasma bilirubin in CCl₄-treated group is possibly because of its leakage from hepatocytes to plasma, generally due to the hepatic obstruction to bile outflow and cholestasis.^[43] In addition, CCl₄ also significantly reduced plasma TP probably causing endoplasmic reticulum destabilization and destruction of protein synthesis.

But the treatment of BEHC associated with Se well-improved plasma levels of TB and TP through stabilizing biliary obstruction and resynthesizing protein. These findings suggest that the combination of BEHC with Se is effective to stabilize CCl₄-induced liver dysfunction in rats.

Oxidative stress dysfunction results from free radicals and reactive oxygen species; these reactive molecules are involved in many physiological processes and human diseases, such as cancer, aging, arthritis, Parkinson's syndrome, ischemia, and liver injury. MDA level is the primary measure of oxidative damage in liver. Toxic radicals as the superoxide radical and also trichloromethyl and trichloromethyl peroxy destabilize cell membranes induced lipid peroxidation.^[44,45] Our data revealed that CCl₄ treatment significantly increased the level of MDA in liver tissues. These findings are in line with a previous

report which revealed that CCl₄ led to an increase in lipid peroxidation.^[35] A significant amelioration of MDA levels was observed in groups treated with BEHC at the dose 400 mg/kg and BEHC associated with Se. These results could be explained by the potential activity of rutin, ferulic acid, and chlorogenic acid that were present in plant extract as found by HPLC analysis. These three components are known for their capacity of quenching free radicals and diminishing lipid peroxidation and therefore acting as potential therapeutic agents.^[46,47]

Antioxidant enzymes such as GSH, CAT, and SOD are considered as the first line of cellular defense against oxidative damage. In this study, the treatment of rats with CCl₄ decreased the antioxidant activity of these enzymes by decreasing their levels in the hepatic tissues of the rat. Our results are similar to other reports that showed that CCl₄ causes a decrease in antioxidant enzymes and an increase in lipid peroxidation.^[19,33] Pretreatment with BEHC at the dose of 400 mg/kg associated with Se ameliorated the deleterious effects of CCl₄ by increasing significantly the levels of oxidative enzymes compared with the normal control group; this antioxidant effect of BEHC was probably related to its capability to reduce the accumulation of free radicals.

The current research is the first one to show the potent antioxidant and protective effect of BEHC in association with Se against oxidative damage *in vivo*.

Conclusion

The present investigation proves that BEHC combined with Se prevented CCl₄-induced liver damage in rats and significantly reduced oxidative stress by increasing antioxidant enzymes activities, decreasing the level of MDA, and preventing the increase in plasma aminotransferases levels. Therefore, due to its higher polyphenol content and potent antioxidant activities, *H. cheirifolia* confirmed its reliability with Se as a hepatoprotective plant to rat against CCl₄-induced liver damage.

Financial support and sponsorship

This study was supported by the Ministry of Higher Education and Scientific Research of Algeria.

Conflicts of interest

There are no conflicts of interest.

References

- Weber LW, Boll M, Stampfl A. Hepatotoxicity and mechanism of action of haloalkanes: Carbon tetrachloride as a toxicological model. *Crit Rev Toxicol* 2003;33:105-36.
- Desai SN, Patel DK, Devkar RV, Patel PV, Ramachandran AV. Hepatoprotective potential of polyphenol rich extract of *Murraya koenigii* L.: An *in vivo* study. *Food Chem Toxicol* 2012;50:310-4.
- Zhou D, Ruan J, Cai Y, Xiong Z, Fu W, Wei A. Antioxidant and hepatoprotective activity of ethanol extract of *Arachniodes exilis* (Hance) Ching. *J Ethnopharmacol* 2010;129:232-7.
- Dzoyem JP, Eloff JN. Anti-inflammatory, anticholinesterase and antioxidant activity of leaf extracts of twelve plants used traditionally to alleviate pain and inflammation in South Africa. *J Ethnopharmacol* 2015;160:194-201.
- Rahmouni F, Hamdaoui L, Badraoui R, Rebai T. Protective effects of *Teucrium polium* aqueous extract and ascorbic acid on hematological and some biochemical parameters against carbon tetrachloride (CCl₄) induced toxicity in rats. *Biomed Pharmacother* 2017;91:43-8.
- Li CC, Hsiang CY, Wu SL, Ho TY. Identification of novel mechanisms of silymarin on the carbon tetrachloride-induced liver fibrosis in mice by nuclear factor-kb bioluminescent imaging-guided transcriptomic analysis. *Food Chem Toxicol* 2012;50:1568-75.
- Rayman MP, Rayman MP. The argument for increasing selenium intake. *Proc Nutr Soc* 2002;61:203-15.
- Akhgar MR, Shariatifar M, Akhgar AR, Moradalizadeh M, Faghihi-Zarandi A. Chemical composition and antibacterial activity of the leaf essential oil from *Hertia intermedia*. *Chem Nat Compd* 2012;48:329-331.
- Kada S, Bouriche H, Senator A, Demirtaş I, Özen T, Çeken Toptancı B, *et al.* Protective activity of *Hertia cheirifolia* extracts against DNA damage, lipid peroxidation and protein oxidation. *Pharm Biol* 2017;55:330-7.
- Segueni N, Zellagui A, Boulechfar S, Derouiche K, Rhouati S. Essential oil of *Hertia cheirifolia* leaves: Chemical composition, antibacterial and antioxidant activities. *J Mtr Env Sci* 2017;8:551-55.
- Ammar S, Edziri H, Mahjoub MA, Chatter R, Bouraoui A, Mighri Z. Spasmolytic and anti-inflammatory effects of constituents from *Hertia cheirifolia*. *Phytomedicine* 2009;16:1156-61.
- Attia S, Grissa KL, Maillieux AC, Heuskin S, Lognay G, Hance T. Acaricidal activities of *Santolina africana* and *Hertia cheirifolia* essential oils against the two-spotted spider mite (*Tetranychus urticae*). *Pest Manag Sci* 2012;68:1069-76.
- Majouli K, Hlila MB, Hamdi A, Flamini G, Jannet HB, Kenani A. Antioxidant activity and α -glucosidase inhibition by essential oils from *Hertia cheirifolia* (L.). *Ind Crops Prod* 2016;82:23-28.
- Cetkovic GS, Djilas SM, Canadanovic-Brunet JM, Tumbas VT. Thin-layer chromatography analysis and scavenging activity of Marigold (*Calendula officinalis* L.) extracts. *Acta Period Technol* 2003;34:1-148.
- Wong CC, Li HB, Cheng KW, Chen F. A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. *Food Chem*. 2006;97:705-711.
- Bahorun T, Gressier B, Trotin F, Brunet C, Dine T, Luyckx M, *et al.* Oxygen species scavenging activity of phenolic extracts from hawthorn fresh plant organs and pharmaceutical preparations. *Arzneimittelforschung* 1996;46:1086-9.
- Barros L, Duenas M, Ferreira ICFR, Baptista P, Santos-Buelga C. Phenolic acids determination by HPLC-DAD-ESI/MS in sixteen different Portuguese wild mushrooms species. *Food Chem Toxicol* 2009;47:1076-9.
- Blois MS. Antioxidant determinations by the use of a stable Free Radical. *Nature* 1958;181:1119-200.
- Mahmoodzadeh Y, Mazani M, Rezagholizadeh L. Hepatoprotective effect of methanolic *Tanacetum parthenium* extract on CCl₄-induced liver damage in rats. *Toxicol Rep* 2017;4:455-62.
- Douhri B, Idaomar M, Skali Senhaji N, Ennabili A, Abrini J. Hepatoprotective Effect of *Origanum elongatum* against carbon tetrachloride (CCl₄) induced toxicity in rats. *Eur J Med Plants* 2014;4:14-28.
- Bouriche H, Kada S, Assaf AM, Senator A, Gül F, Dimertas I. Phytochemical screening and anti-inflammatory properties of Algerian *Hertia cheirifolia* methanol extract. *Pharm Biol* 2016;54:2584-90.
- Iqbal M, Sharma SD, Okazaki Y, Fujisawa M, Okada S. Dietary supplementation of curcumin enhances antioxidant and phase II

- metabolizing enzymes in ddY male mice: Possible role in protection against chemical carcinogenesis and toxicity. *Pharmacol Toxicol* 2003;92:33-8.
23. Claiborne A. Catalase activity. In: Greenwald RA, editor. *CRC Handbook of Methods in Oxygen Radical Research*. Boca Raton, FL: CRS Press; 1985.
 24. Beauchamp C, Fridovich I. Assay of superoxide dismutase. *Anal Biochem* 1971;44:276-87.
 25. Ellman GL. Plasma antioxidants. *Arch Biochem Biophys* 1959;82:70-7.
 26. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351-8.
 27. Ibrahim NA, Mohammed MMD, Aly HF, Ali SA, Al-Hady DA. Efficiency of the leaves and fruits of *Aegle marmelos* methanol extract (L.) Correa and their relative hepatotoxicity induced by CCl₄ and identification of their active constituents by using LC/MS/MS. *Toxicol Rep* 2018;5:1161-8.
 28. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol* 2011;48:412-22.
 29. Majouli K, Mahjoub MA, Rahim F, Hamdi A, Wadood A, Besbes Hlila M, *et al.* Biological properties of *Hertia cheirifolia* L. Flower extracts and effect of the nopol on α -glucosidase. *Int J Biol Macromol* 2017;95:757-61.
 30. Majouli K, Hamdi A, Hlila MB. Phytochemical analysis and biological activities of *Hertia cheirifolia* L. Roots extracts. *Asian Pac J Trop Med* 2017;10:1134-9.
 31. Menakh M, Mahdi D, Boutellaa S, Zellagui A, Lahouel M, Bensouici C. *In vitro* antioxidant activity and protective effect of *Hertia cheirifolia* L. n-butanol extract against liver and heart mitochondrial oxidative stress in rat. *Acta Sci Nat* 2020;7:33-45.
 32. Zielinska D, Szawara-Nowak D, Zieliński H. Determination of the antioxidant activity of rutin and its contribution to the antioxidant capacity of diversified buckwheat origin material by updated analytical strategies. *Pol J Food Nutr Sci* 2010;60:315-21.
 33. Allis JW, Ward TR, Seely JC, Simmons JE. Assessment of hepatic indicators of subchronic carbon tetrachloride injury and recovery in rats. *Fundam Appl Toxicol* 1990;15:558-70.
 34. Klaassen CD, Watkins JB 3rd. Mechanisms of bile formation, hepatic uptake, and biliary excretion. *Pharmacol Rev* 1984;36:1-67.
 35. Hsouna AB, Saoudi M, Trigui M, Jamoussi K, Boudawara T, Jaoua S, *et al.* Characterization of bioactive compounds and ameliorative effects of *Ceratonia siliqua* leaf extract against CCl₄ induced hepatic oxidative damage and renal failure in rats. *Food Chem Toxicol* 2011;49:3183-91.
 36. Majouli K, Hamdi A, Abdelhamid A, Bouraoui A, Kenani A. Anti-inflammatory activity and gastroprotective effect of *Hertia cheirifolia* L. Roots extract. *J Ethnopharmacol* 2018;217:7-10.
 37. Mézes M, Balogh K. Prooxidant mechanisms of selenium toxicity: A review. *Acta Biol Szeged* 2009;53:15-18.
 38. Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 2006;160:1-40.
 39. Ding M, Potter JJ, Liu X, Torbenson MS, Mezey E. Selenium supplementation decreases hepatic fibrosis in mice after chronic carbon tetrachloride administration. *Biol Trace Elem Res* 2010;133:83-97.
 40. Fernandez ML, West KL. Mechanisms by which dietary fatty acids modulate plasma lipids. *J Nutr* 2005;135:2075-8.
 41. Marimuthu S, Adluri RS, Rajagopalan R, Menon VP. Protective role of ferulic acid on carbon tetrachloride-induced hyperlipidemia and histological alterations in experimental rats. *J Basic Clin Physiol Pharmacol* 2013;24:59-66.
 42. Klyszejko B, Lyzzywek G. Effects of a sublethal concentration of deltamethrin on biochemical parameter of blood serum of carp (*Cyprinus carpio* L). *Acta Ichtyolo Piscat* 1999;292:109-116.
 43. Field KM, Dow C, Michael M. Part I: Liver function in oncology: Biochemistry and beyond. *Lancet Oncol* 2008;9:1092-101.
 44. Klibet F, Boumendjel A, Abdenour C, Bouzerna N, Boulakoud MS, El Feki A. Hepatoprotective role and antioxidant capacity of selenium on arsenic-induced liver injury in rats. *Exper Toxicol Path* 2012;64:167-74.
 45. Senosy W, Kamal A, El-Toumy S, El-Gendy EH. Phenolic compounds and hepatoprotective activity of *Centaurea aegyptiaca* L. on carbon tetrachloride-induced hepatotoxicity in rats. *J Adv Pharm Res* 2018;2:123-32.
 46. Yan JJ, Cho JY, Kim HS, Kim KL, Jung YS, Huh SO, *et al.* Protection against β -amyloid peptide toxicity *in vivo* with long-term administration of ferulic acid. *British J Pharma* 2001;133:89-96.
 47. Naveed M, Hejazi V, Abbas M, Kamboh AA, Khan GJ, Shumzaid M, *et al.* Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed Pharmacother* 2018;97:67-74.