



ARTICLE

Modeling the Spread of Tuberculosis with Piecewise Differential Operators

Abdon Atangana^{1,2} and Ilknur Koca^{3,*}

¹Institute for Groundwater Studies, Faculty of Natural and Agricultural Sciences, University of the Free State, Bloemfontein, 9301, South Africa

²Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, 008864, Taiwan

³Department of Accounting and Financial Management, Seydikemer High School of Applied Sciences, Mugla Sitki Kocman University, Mugla, 48300, Turkey

*Corresponding Author: Ilknur Koca. Email: ilknurkoca@mu.edu.tr

Received: 09 September 2021 Accepted: 09 November 2021

ABSTRACT

Very recently, a new concept was introduced to capture crossover behaviors that exhibit changes in patterns. The aim was to model real-world problems exhibiting crossover from one process to another, for example, randomness to a power law. The concept was called piecewise calculus, as differential and integral operators are defined piecewisely. These behaviors have been observed in the spread of several infectious diseases, for example, tuberculosis. Therefore, in this paper, we aim at modeling the spread of tuberculosis using the concept of piecewise modeling. Several cases are considered, conditions under which the unique system solution is obtained are presented in detail. Numerical simulations are performed with different values of fractional orders and density of randomness.

KEYWORDS

Spread of tuberculosis; piecewise differentiation; numerical simulation

1 Introduction

Although several studies have been done on behaviors of the tuberculosis virus, its spread, and its effect on the human's body until today, this virus persists and kills humans around the world each year. It is even believed that the tuberculosis virus has affected about 25 percent of the world population since about one percent of the world population is infected each year according to what is reported in the literatures [1–4]. Tuberculosis is a seasonal transmissible disease, as the peaks are reached every spring and summer. However, there is no apparent scientific reason recorded that can explain this variation. Nevertheless, it is recorded that the virus spreads more during weather conditions like low temperature, humidity, and low rainfall. Thus tuberculosis incidence rates could be linked to change in the climate. Having peaks that occurred during some period of the year show that the spread had many waves since antiquity. Indeed, each wave has a specific pattern different from others or similar in some cases. It can be concluded that the virus spread follows piecewise patterns. Mathematicians have tried to provide mathematical models to



depict the spread behaviors as a function of time. Several studies have been performed in the decades. The reproductive number of this virus has been calculated in many studies. New and modified models have been provided and studied in detail. Several differential and integral operators have been used, for example, fractional differential operators to replicate spread behaviors. Fractional derivative based on power law was introduced to replicate behaviors resembling the power law [5–11]. Different techniques have been employed, for example, the stochastic process to capture random behaviors. Nevertheless, the problem of different was not really addressed. The concept of piecewise differential and integral operators was recently suggested and employed to model some complex real-world problems, such as chaos and other epidemiological problems [12]. The concept seems to be efficient when modeling problems with crossover behaviors. In this paper, we aim to modify an existing tuberculosis model with the concept of piecewise differentiation.

1.1 Important Definitions of Fractional Modelling

Definition 1: Let $\alpha > 0$ of a function $h : (0, \infty) \rightarrow R$ and the Riemann-Liouville derivative of fractional order is presented as

$$D_t^\alpha h(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-x)^{-\alpha} h(x) dx, \quad 0 < \alpha \leq 1. \quad (1)$$

Definition 2: Let $h : H^1(a, b)$, $b > a$, $0 < \alpha < 1$ then, the Caputo-Fabrizio derivative of fractional derivative is presented as

$${}^a_{CF}D_t^\alpha h(t) = \frac{1}{1-\alpha} \int_a^t h'(x) \exp\left[-\alpha \frac{(t-x)}{1-\alpha}\right] dx. \quad (2)$$

Definition 3: Let $h : H^1(a, b)$, $b > a$, $\alpha \in (0, 1)$ then, the definition of the new fractional derivative (Atangana-Baleanu derivative in Caputo sense) is presented as

$${}^a_{ABC}D_t^\alpha h(t) = \frac{AB(\alpha)}{1-\alpha} \int_a^t h'(x) E_\alpha\left[-\alpha \frac{(t-x)^\alpha}{1-\alpha}\right] dx, \quad (3)$$

where ${}^a_{ABC}D_t^\alpha$ is fractional operator with Mittag-Leffler kernel in the Caputo sense with order α with respect to t and

$$AB(\alpha) = 1 - \alpha + \frac{\alpha}{\Gamma(\alpha)}, \quad (4)$$

is a normalization function.

Definition 4: Let h be continuous not necessary differentiable in $[t_1, T]$. Thus, the piecewise Riemann-Liouville derivative is presented as

$${}^0_{PRL}D_t^\alpha h(t) = \begin{cases} h'(t), & \text{if } 0 \leq t \leq t_1 \\ {}^{RL}_{t_1}D_t^\alpha h(t), & \text{if } t_1 \leq t \leq T \end{cases}, \quad (5)$$

where ${}^0_{PRL}D_t^\alpha$ presents classical derivative on $0 \leq t \leq t_1$ and Riemann-Liouville fractional derivative on $t_1 \leq t \leq T$.

Definition 5: The piecewise derivative with classical and exponential decay kernel is defined as

$${}^{\text{PCF}}_0 D_t^\alpha h(t) = \begin{cases} h'(t), & \text{if } 0 \leq t \leq t_1 \\ {}^{\text{CF}}_{t_1} D_t^\alpha h(t), & \text{if } t_1 \leq t \leq T \end{cases} \quad (6)$$

and

$${}^{\text{PCF}}_0 D_t^\alpha h(t) = \begin{cases} h'(t), & \text{if } 0 \leq t \leq t_1 \\ {}^{\text{CFR}}_{t_1} D_t^\alpha h(t), & \text{if } t_1 \leq t \leq T \end{cases} \quad (7)$$

where ${}^{\text{PCF}}_0 D_t^\alpha$ presents classical derivative on $0 \leq t \leq t_1$ and Caputo-Fabrizio fractional derivative on $t_1 \leq t \leq T$.

Definition 6: The piecewise derivative with classical and Mittag-Leffler kernel is given as

$${}^{\text{PAB}}_0 D_t^\alpha h(t) = \begin{cases} h'(t), & \text{if } 0 \leq t \leq t_1 \\ {}^{\text{ABC}}_{t_1} D_t^\alpha h(t), & \text{if } t_1 \leq t \leq T \end{cases} \quad (8)$$

where ${}^{\text{PAB}}_0 D_t^\alpha$ presents classical derivative on $0 \leq t \leq t_1$ and Atangana-Baleanu fractional derivative on $t_1 \leq t \leq T$.

Definition 7: Let h be continuous and $\alpha > 0$ then a piecewise integral of h is given as

$${}^{\text{PPL}}_0 J_t^\alpha h(t) = \begin{cases} \int_0^{t_1} h(\tau) d\tau, & \text{if } 0 \leq t \leq t_1 \\ \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-\tau)^{\alpha-1} h(\tau) d\tau, & \text{if } t_1 \leq t \leq T \end{cases} \quad (9)$$

where ${}^{\text{PPL}}_0 J_t^\alpha h(t)$ presents classical integral on $0 \leq t \leq t_1$ and the integral with power-law kernel on $t_1 \leq t \leq T$.

Definition 8: Let h be continuous and $\alpha > 0$ then a piecewise integral of h is given as

$${}^{\text{PCF}}_0 J_t^\alpha h(t) = \begin{cases} \int_0^{t_1} h(\tau) d\tau, & \text{if } 0 \leq t \leq t_1 \\ \frac{1-\alpha}{M(\alpha)} h(t) + \frac{\alpha}{M(\alpha)} \int_{t_1}^t h(\tau) d\tau, & \text{if } t_1 \leq t \leq T \end{cases} \quad (10)$$

where ${}^{\text{PCF}}_0 J_t^\alpha h(t)$ presents classical integral on $0 \leq t \leq t_1$ and Caputo-Fabrizio integral on $t_1 \leq t \leq T$.

2 Tuberculosis Epidemic Model

In this section, we take into account the following piecewise model of tuberculosis:

$$\begin{aligned}
 \frac{dS(t)}{dt} &= \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\
 \frac{dE(t)}{dt} &= \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\
 \frac{dI_1(t)}{dt} &= p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\
 \frac{dI_2(t)}{dt} &= \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t).
 \end{aligned} \tag{11}$$

The initial conditions are taken as follows:

$$S(0) = S_0, \quad E(0) = E_0, \quad I_1(0) = A_0, \quad I_2(0) = I_0. \tag{12}$$

But we noted that the model was considered with its classical version in paper [13] before. Now, we give the meanings of the parameters of model considered in this paper given by Table 1 below:

Table 1: The meanings of the parameters of model

S:	→ susceptible individuals
E:	→ exposed individuals
I_1 :	→ first infected class
I_2 :	→ second infected class
λ :	→ the recruitment rate
β_1 :	→ the level of contact with infectious I_1
β_2 :	→ the level of contact with infectious I_1
μ :	→ the rate of natural death
δ :	→ death rate from disease in the TB infected individuals
γ :	→ moving an individual from the latent sub-population to the infected sub-population
ϕr_1 :	→ first line treatment
r :	→ $(1 - r_1)$, $0 < r_1 < 1$
$\phi(1 - r_1)$:	→ the fraction of the infectious class
p_1 :	→ $(1 - p)$, $0 < p < 1$
q_1 :	→ $(1 - q)$, $0 < q < 1$

2.1 Second Derivative of Lyapunov Function and Strength Number

Lyapunov function formulation has been used in different analyses in different fields in the last past year. In epidemiology, this function has been used to determine the stability analysis of an epidemiological model. It has been reported that the Lyapunov can be viewed as energy; therefore, a sign of the first derivative of the function can be useful for the determination of stability. Nevertheless, the sign of the first derivative of a function may not be enough to define whether we have a local maximum or local minimum. On this note, it was suggested that the sign

of the second derivative should also be studied. In this section, we shall proceed with such analysis to determine the sign of our model's second derivative of the associate Lyapunov function.

In this section, we present the second derivative of Lyapunov function for the model [2–14].

$$\begin{aligned}
\frac{dS(t)}{dt} &= \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\
\frac{dE(t)}{dt} &= \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\
\frac{dI_1(t)}{dt} &= p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\
\frac{dI_2(t)}{dt} &= \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t).
\end{aligned} \tag{13}$$

Now we find second derivative of Lyapunov function for model with following equality:

$$\begin{aligned}
\ddot{L} &= \frac{d\dot{L}}{dt} = \frac{d}{dt} \left\{ \left(1 - \frac{S^*}{S}\right) \dot{S} + \left(1 - \frac{E^*}{E}\right) \dot{E} \right. \\
&\quad \left. + \left(1 - \frac{I_1^*}{I_1}\right) \dot{I}_1 + \left(1 - \frac{I_2(t)^*}{I_2(t)}\right) \dot{I}_2 \right\}, \\
&= \left(\frac{\dot{S}}{S}\right)^2 S^* + \left(\frac{\dot{E}}{E}\right)^2 E^* + \left(\frac{\dot{I}_1}{I_1}\right)^2 I_1^* + \left(\frac{\dot{I}_2}{I_2}\right)^2 I_2^* + \left(1 - \frac{S^*}{S}\right) \ddot{S} + \left(1 - \frac{E^*}{E}\right) \ddot{E} \\
&\quad + \left(1 - \frac{I_1^*}{I_1}\right) \ddot{I}_1 + \left(1 - \frac{I_2^*}{I_2}\right) \ddot{I}_2.
\end{aligned} \tag{14}$$

Here second derivatives of classes are given as below:

$$\begin{aligned}
\ddot{S}(t) &= -\beta_1 (\dot{S}(t)I_1(t) + \dot{I}_1(t)S(t)) - \beta_2 (\dot{S}(t)I_2(t) + \dot{I}_2(t)S(t)) - \mu \dot{S}(t), \\
\ddot{E}(t) &= \beta_1 p_1 (\dot{S}(t)I_1(t) + \dot{I}_1(t)S(t)) + \beta_2 q_1 (\dot{S}(t)I_2(t) + \dot{I}_2(t)S(t)) - (\mu + \gamma) \dot{E}(t), \\
\ddot{I}_1(t) &= \beta_1 p (\dot{S}(t)I_1(t) + \dot{I}_1(t)S(t)) + \beta_2 q (\dot{S}(t)I_2(t) + \dot{I}_2(t)S(t)) + \gamma \dot{E}(t) - (\phi + \mu + \delta_1) \dot{I}_1(t), \\
\ddot{I}_2(t) &= \phi(1 - r_1) \dot{I}_1(t) - (\mu + \delta_2) \dot{I}_2(t) - \varphi r_2 \dot{I}_2(t).
\end{aligned} \tag{15}$$

Then we have,

$$\begin{aligned}
\frac{d\dot{L}}{dt} &= \left(\frac{\dot{S}}{S}\right)^2 S^* + \left(\frac{\dot{E}}{E}\right)^2 E^* + \left(\frac{\dot{I}_1}{I_1}\right)^2 I_1^* + \left(\frac{\dot{I}_2}{I_2}\right)^2 I_2^* \\
&\quad + \left(1 - \frac{S^*}{S}\right) \left\{ \begin{array}{l} -\beta_1 (\dot{S}(t)I_1(t) + \dot{I}_1(t)S(t)) \\ -\beta_2 (\dot{S}(t)I_2(t) + \dot{I}_2(t)S(t)) - \mu \dot{S}(t) \end{array} \right\}
\end{aligned} \tag{16}$$

$$\begin{aligned}
& + \left(1 - \frac{E^*}{E}\right) \left\{ \beta_1 p_1 \left(\dot{S}(t) I_1(t) + \dot{I}_1(t) S(t) \right) \right. \\
& \quad \left. + \beta_2 q_1 \left(\dot{S}(t) I_2(t) + \dot{I}_2(t) S(t) \right) - (\mu + \gamma) \dot{E}(t) \right\} \\
& + \left(1 - \frac{I_1^*}{I_1}\right) \left\{ \beta_1 p \left(\dot{S}(t) I_1(t) + \dot{I}_1(t) S(t) \right) \right. \\
& \quad \left. + \beta_2 q \left(\dot{S}(t) I_2(t) + \dot{I}_2(t) S(t) \right) \right. \\
& \quad \left. + \gamma \dot{E}(t) - (\phi + \mu + \delta_1) \dot{I}_1(t) \right\} \\
& + \left(1 - \frac{I_2^*}{I_2}\right) \left\{ \phi(1 - r_1) \dot{I}_1(t) - (\mu + \delta_2) \dot{I}_2(t) - \varphi r_2 \dot{I}_2(t) \right\}.
\end{aligned}$$

$$\frac{d\dot{L}}{dt} = \dot{L}(S, E, I_1, I_2) \tag{17}$$

$$\begin{aligned}
& + \dot{S}(t) \left\{ \begin{array}{l} -\beta_1 I_1(t) - \beta_2 I_2(t) - \mu - \frac{E^*}{E} \beta_1 p_1 I_1(t) \\ -\frac{E^*}{E} \beta_2 q_1 I_2(t) - I_1^* \beta_1 p - \frac{I_1^*}{I_1} \beta_2 q I_2(t) \\ +\beta_1 p_1 I_1(t) + \beta_2 q_1 I_2(t) + \beta_1 p I_1(t) \\ +\beta_2 q I_2(t) + \frac{S^*}{S} \beta_1 I_1(t) + \frac{S^*}{S} \beta_2 I_2(t) + \frac{S^*}{S} \mu \end{array} \right\} \\
& + \dot{E}(t) \left\{ -(\mu + \gamma) - \frac{I_1^*}{I_1} \gamma + \gamma \right\} \\
& + \dot{I}_1(t) \left\{ \begin{array}{l} -\beta_1 S(t) - (\phi + \mu + \delta_1) - \frac{E^*}{E} \beta_1 p_1 S(t) \\ -\frac{I_1^*}{I_1} \beta_1 p S(t) - \frac{I_2^*}{I_2} \phi(1 - r_1) + \beta_1 p_1 S(t) \\ +\beta_1 p S(t) + \phi(1 - r_1) + S^* \beta_1 + \frac{I_1^*}{I_1} (\phi + \mu + \delta_1) \end{array} \right\} \\
& + \dot{I}_2(t) \left\{ \begin{array}{l} -\beta_2 S(t) - (\mu + \delta_2) - \varphi r_2 \\ -\frac{E^*}{E} \beta_2 q_1 S(t) - \frac{I_1^*}{I_1} \beta_2 q S(t) + \beta_2 q_1 S(t) \\ +\beta_2 q S(t) + S^* \beta_2 + \frac{I_2^*}{I_2} (\mu + \delta_2) + \frac{I_2^*}{I_2} \varphi r_2 \end{array} \right\}.
\end{aligned}$$

Now replacing $\dot{S}(t)$, $\dot{E}(t)$, $\dot{I}_1(t)$, and $\dot{I}_2(t)$ by their respective formula with their positive and negative parts, we have

$$\frac{d^2 L}{dt^2} = \dot{L}(S, E, I_1, I_2) + \Pi^+ - \Pi^-, \tag{18}$$

$$\frac{d^2 L}{dt^2} = \underbrace{\dot{L}(S, E, I_1, I_2) + \Pi^+}_{\Pi_1} - \underbrace{\Pi^-}_{\Pi_2}$$

where

$$\begin{aligned}
\Pi^+ = & \lambda \left(\begin{array}{l} \beta_1 p_1 I_1(t) + \beta_2 q_1 I_2(t) + \beta_1 p I_1(t) \\ + \beta_2 q I_2(t) + \frac{S^*}{S} \beta_1 I_1(t) + \frac{S^*}{S} \beta_2 I_2(t) + \frac{S^*}{S} \mu \end{array} \right) + (\beta_1 S(t) I_1(t) + \beta_2 S(t) I_2(t) + \mu S(t)) \\
& \cdot \left(\begin{array}{l} \beta_1 I_1(t) + \beta_2 I_2(t) + \mu + \frac{E^*}{E} \beta_1 p_1 I_1(t) \\ + \frac{E^*}{E} \beta_2 q_1 I_2(t) + I_1^* \beta_1 p + \frac{I_1^*}{I_1} \beta_2 q I_2(t) \end{array} \right) + (\beta_1 p_1 S(t) I_1(t) + \beta_2 q_1 S(t) I_2(t)) \gamma \\
& + ((\mu + \gamma) E(t)) \left((\mu + \gamma) + \frac{I_1^*}{I_1} \right) \\
& + \left(\begin{array}{l} \beta_1 p I_1(t) S(t) + \beta_2 q I_2(t) S(t) \\ + \gamma E(t) \end{array} \right) \left(\begin{array}{l} \beta_1 p_1 S(t) + \beta_1 p S(t) \\ + \phi(1 - r_1) + S^* \beta_1 \\ + \frac{I_1^*}{I_1} ((\phi + \mu + \delta_1)) \end{array} \right) \\
& + (\phi + \mu + \delta_1) I_1(t) \left(\begin{array}{l} \beta_1 S(t) + (\phi + \mu + \delta_1) \\ + \frac{E^*}{E} \beta_1 p_1 S(t) + \frac{I_1^*}{I_1} \beta_1 p S(t) + \frac{I_2^*}{I_2} \phi(1 - r_1) \end{array} \right) \\
& + \phi(1 - r_1) I_1(t) \left(\begin{array}{l} \beta_2 q_1 S(t) + \beta_2 q S(t) + S^* \beta_2 \\ + \frac{I_2^*}{I_2} (\mu + \delta_2) + \frac{I_2^*}{I_2} \varphi r_2 \\ + (\varphi r_2 I_2) \left(\begin{array}{l} \beta_2 S(t) + (\mu + \delta_2) + \varphi r_2 \\ + \frac{E^*}{E} \beta_2 q_1 S(t) + \frac{I_1^*}{I_1} \beta_2 q S(t) \end{array} \right) \end{array} \right). \tag{19} \\
\Pi^- = & \lambda \left(\begin{array}{l} \beta_1 I_1(t) + \beta_2 I_2(t) + \mu + \frac{E^*}{E} \beta_1 p_1 I_1(t) \\ + \frac{E^*}{E} \beta_2 q_1 I_2(t) + \beta_1 p I_1^*(t) + \frac{I_1^*}{I_1} \beta_2 q I_2(t) \end{array} \right) + (\beta_1 S(t) I_1(t) + \beta_2 S(t) I_2(t) + \mu S(t)) \\
& \cdot \left(\begin{array}{l} \beta_1 p_1 I_1(t) + \beta_2 q_1 I_2(t) + \beta_1 p I_1(t) + \beta_2 q I_2(t) \\ \frac{S^*}{S} \beta_1 I_1(t) + \frac{S^*}{S} \beta_2 I_2(t) + \frac{S^*}{S} \mu \end{array} \right) \\
& + (\beta_1 p_1 S(t) I_1(t) + \beta_2 q_1 S(t) I_2(t)) \left(-(\mu + \gamma) - \frac{I_1^*}{I_1} \gamma \right) - \gamma (\mu + \gamma) E(t) \\
& + (\phi + \mu + \delta_1) I_1(t) \left(\begin{array}{l} \beta_1 p_1 S(t) + \frac{I_1^*}{I_1} (\phi + \mu + \delta_1) \\ + \beta_1 p S(t) + \beta_1 S^* + \phi(1 - r_1) \end{array} \right) \\
& + (\beta_1 p S(t) I_1(t) + \beta_2 q S(t) I_2(t) + \gamma E(t))
\end{aligned}$$

$$\begin{aligned} & \left(\begin{array}{l} \beta_1 S(t) + (\phi + \mu + \delta_1) + \frac{E^*}{E} \beta_1 p_1 S(t) \\ + \frac{I_1^*}{I_1} \beta_1 p S(t) + \frac{I_2^*}{I_2} \phi (1 - r_1) \end{array} \right) \\ & + \phi (1 - r_1) I_1(t) \left(\begin{array}{l} \beta_2 S(t) + (\mu + \delta_2) + \varphi r_2 \\ + \frac{E^*}{E} \beta_2 q_1 S(t) + \frac{I_1^*}{I_1} \beta_2 q S(t) \\ + (\varphi r_2 I_2) \left(\begin{array}{l} \beta_2 q_1 S(t) + q \beta_2 S(t) + S^* \beta_2 \\ + \frac{I_2^*}{I_2} (\mu + \delta_2) + \frac{I_2^*}{I_2} \varphi r_2 \end{array} \right) \end{array} \right). \end{aligned}$$

Now we can easily put following results for obtained results above:

$$\frac{d^2 L}{dt^2} = \Pi_1 - \Pi_2. \quad (20)$$

Then

$$\text{If } \Pi_1 > \Pi_2 \text{ then } \frac{d^2 L}{dt^2} > 0, \quad (21)$$

$$\text{If } \Pi_1 < \Pi_2 \text{ then } \frac{d^2 L}{dt^2} < 0,$$

$$\text{If } \Pi_1 = \Pi_2 \text{ then } \frac{d^2 L}{dt^2} = 0.$$

So, the interpretation associated the sign of second order.

2.2 Strength Number

Without a doubt, the reproductive number has been utilized as a powerful mathematical tool to the stability of a mathematical model for a given infectious disease. While it has been used with some success, it has also been criticized as an insufficient tool to predict the behavior of the spread. For example, it was pointed out that there are several ways to obtain this value on the other hand. However, it was also argued that this value could not help humans to determine whether the model will determine waves. The concept of strength number has been suggested to further the analysis and will be used in this section.

The component F_A is obtained with deriving the nonlinear part of the infected classes. In our model there are two infected classes named by I_1 and I_2 . These infected classes given by

$$\dot{I}_1 = p \beta_1 S(t) I_1(t) + q \beta_2 S(t) I_2(t) \quad (22)$$

$$+ \gamma E(t) - (\phi + \mu + \delta_1) I_1(t),$$

$$\dot{I}_2 = \phi (1 - r_1) I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t).$$

But we only use nonlinear part of infected classes. So we use \dot{I}_2 classes. Nonlinear part of \dot{I}_1 classes is given by

$$= S(t)I_1(t) + S(t)I_2(t), \quad (23)$$

$$\frac{\partial}{\partial I_1} (S(t)I_1(t) + S(t)I_2(t)) = S(t), \quad (24)$$

$$\frac{\partial^2}{\partial I_1^2} (S(t)) = 0.$$

In this case, we can have the following

$$F_A = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \quad (25)$$

Then

$$\det(F_A V^{-1} - \lambda I) = 0, \quad (26)$$

leads to

$$A_0 = 0. \quad (27)$$

A_0 means there is no strength. Also there are more conclusion when strength is zero

- 1) The disease will spread with a constant speed.
- 2) The disease will not renewal process therefore no new wave will be expected.
- 3) The magnitude of the spread will be the same at all time until extinction.

3 Applications of Piecewise Derivative

3.1 A Mathematical Model of Tuberculosis Epidemic Model with Piecewise Modeling

In this section, we present some applications of piecewise derivative for tuberculosis epidemic model such as

$$\begin{cases} \frac{dS(t)}{dt} = \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\ \frac{dE(t)}{dt} = \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\ \frac{dI_1(t)}{dt} = p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\ \frac{dI_2(t)}{dt} = \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t). \end{cases} \quad \text{if } 0 \leq t \leq W_1, \quad (28)$$

$$\begin{cases} {}^C D_t^\alpha S(t) = \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\ {}^C D_t^\alpha E(t) = \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\ {}^C D_t^\alpha I_1(t) = p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\ {}^C D_t^\alpha I_2(t) = \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t). \end{cases} \quad \text{if } W_1 \leq t \leq W_2, \quad (29)$$

$$\begin{cases} dS(t) = [\lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t)] dt + \sigma_1 S dB_1(t), \\ dE(t) = \begin{bmatrix} \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) \\ -(\mu + \gamma) E(t) \end{bmatrix} dt + \sigma_2 E dB_2(t), \\ dI_1(t) = \begin{bmatrix} p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) \\ -(\phi + \mu + \delta_1) I_1(t) \end{bmatrix} dt + \sigma_3 I_1 dB_3(t), \\ dI_2(t) = \begin{bmatrix} \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) \\ -\varphi r_2 I_2(t) \end{bmatrix} dt + \sigma_4 I_2 dB_4(t). \end{cases} \quad \text{if } W_2 \leq t \leq W. \quad (30)$$

Let us give necessary conditions for the existence and uniqueness, we must prove that $\forall [0, W_1]$ and $[W_1, W_2]$ $f_i(S, E, I_1, I_2)$ for $i = 1, 2, 3, 4$ satisfy

1) Linear growth condition

$$|f_i(x_i, t)|^2 \leq k_i(1 + |x_i|^2) \quad \text{for } i = 1, 2, 3, 4. \quad (31)$$

and

2) The Lipschitz condition.

$$|f_i(x_i^1, t) - f_i(x_i^2, t)|^2 \leq \bar{k}_i |x_i^1 - x_i^2|^2 \quad \text{for } i = 1, 2, 3, 4. \quad (32)$$

Now we define the norm $\|\varphi\|_\infty = \sup_{t \in D_\varphi} |\varphi(t)|$. Now we put forth the existence and uniqueness of the solution piecewisely for $[0, W_2]$. For $[0, W_2]$, there exist 4 positive constant M_1, M_2, M_3 and $M_4 < \infty$ such that

$$\|S\|_\infty < M_1, \quad (33)$$

$$\|E\|_\infty < M_2,$$

$$\|I_1\|_\infty < M_3,$$

$$\|I_2\|_\infty < M_4.$$

Let us write system as below:

$$\begin{cases} \dot{S} = f_1(S, E, I_1, I_2), \\ \dot{E} = f_2(S, E, I_1, I_2), \\ \dot{I}_1 = f_3(S, E, I_1, I_2), \\ \dot{I}_2 = f_4(S, E, I_1, I_2). \end{cases} \quad \text{if } 0 \leq t \leq W_2. \quad (34)$$

For proof, we consider the function

$$\begin{aligned}
 |f_1(S, E, I_1, I_2)|^2 &= |\lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t)|^2, \\
 &\leq 4\lambda^2 + 4|\beta_1 S(t)I_1(t)|^2 + 4|\beta_2 S(t)I_2(t)|^2 + 4|\mu S(t)|^2, \\
 &\leq 4\lambda^2 + 4\|\beta_1 I_1(t)\|^2 |S(t)|^2 \\
 &\quad + 4\|\beta_2 I_2(t)\|^2 |S(t)|^2 + 4\mu^2 |S(t)|^2, \\
 &\leq 4\lambda^2 + 4\left(\|\beta_1 I_1(t)\|^2 + \|\beta_2 I_2(t)\|^2 + \mu^2\right) |S(t)|^2 \\
 &\leq 4\lambda^2 \left(1 + \frac{\|\beta_1 I_1(t)\|^2 + \|\beta_2 I_2(t)\|^2 + \mu^2}{\lambda^2} |S(t)|^2\right).
 \end{aligned} \tag{35}$$

under the condition that

$$\frac{\|\beta_1 I_1(t)\|^2 + \|\beta_2 I_2(t)\|^2 + \mu^2}{\lambda^2} < 1, \tag{36}$$

then we have

$$|f_1(S, E, I_1, I_2)|^2 \leq k_1(1 + |S(t)|^2). \tag{37}$$

Using same routine

$$\begin{aligned}
 |f_2(S, E, I_1, I_2)|^2 &= |\beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t)|^2, \\
 &\leq 3|\beta_1 p_1 S(t)I_1(t)|^2 + 3|\beta_2 q_1 S(t)I_2(t)|^2 + 3|(\mu + \gamma) E(t)|^2, \\
 &\leq 3 \sup_{t \in [0, T_2]} |\beta_1 p_1 S(t)I_1(t)|^2 \\
 &\quad + 3 \sup_{t \in [0, T_2]} |\beta_2 q_1 S(t)I_2(t)|^2 + 3|(\mu + \gamma) E(t)|^2, \\
 &\leq 3\left\|(\beta_1 p_1 S(t)I_1(t))^2\right\|_\infty \\
 &\quad + 3\left\|(\beta_2 q_1 S(t)I_2(t))^2\right\|_\infty + 3(\mu + \gamma)^2 |E(t)|^2, \\
 &\leq 3\left\|(\beta_1 p_1 S(t)I_1(t))^2\right\|_\infty + 3\left\|(\beta_2 q_1 S(t)I_2(t))^2\right\|_\infty \\
 &\quad \cdot \left(1 + \frac{3(\mu + \gamma)^2}{3\left\|(\beta_1 p_1 S(t)I_1(t))^2\right\|_\infty + 3\left\|(\beta_2 q_1 S(t)I_2(t))^2\right\|_\infty} |E(t)|^2\right),
 \end{aligned} \tag{38}$$

under the condition

$$\frac{(\mu + \gamma)^2}{\left\|(\beta_1 p_1 S(t) I_1(t))^2\right\|_\infty + \left\|(\beta_2 q_1 S(t) I_2(t))^2\right\|_\infty} < 1, \text{ then} \quad (39)$$

$$|f_2(S, E, I_1, I_2)|^2 \leq k_2(1 + |E(t)|^2). \quad (40)$$

For the function $f_3(S, E, I_1, I_2)$

$$\begin{aligned} |f_3(S, E, I_1, I_2)|^2 &= |p\beta_1 S(t) I_1(t) + q\beta_2 S(t) I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t)|^2, \quad (41) \\ &\leq 4 |p\beta_1 S(t) I_1(t)|^2 + 4 |q\beta_2 S(t) I_2(t)|^2 \\ &\quad + 4 |\gamma E(t)|^2 + 4 |(\phi + \mu + \delta_1) I_1(t)|^2, \\ &\leq 4 |p\beta_1 S(t)|^2 |I_1(t)|^2 + 4 |q\beta_2 S(t) I_2(t)|^2 \\ &\quad + 4 |\gamma E(t)|^2 + 4 (\phi + \mu + \delta_1)^2 |I_1(t)|^2, \\ &\leq 4 \sup_{t \in [0, T_2]} |p\beta_1 S(t)|^2 |I_1(t)|^2 + 4 \sup_{t \in [0, T_2]} |q\beta_2 S(t) I_2(t)|^2 \\ &\quad + 4 \sup_{t \in [0, T_2]} |\gamma E(t)|^2 + 4 (\phi + \mu + \delta_1)^2 |I_1(t)|^2, \\ &\leq 4 \left\| (p\beta_1 S(t))^2 \right\|_\infty |I_1(t)|^2 + 4 \left\| (q\beta_2 S(t) I_2(t))^2 \right\|_\infty \\ &\quad + 4 \left\| (\gamma E(t))^2 \right\|_\infty + 4 (\phi + \mu + \delta_1)^2 |I_1(t)|^2, \\ &\leq 4 \left\| (q\beta_2 S(t) I_2(t))^2 \right\|_\infty \\ &\quad + 4 \left\| (\gamma E(t))^2 \right\|_\infty \left(1 + \frac{\left\| (p\beta_1 S(t))^2 \right\|_\infty + (\phi + \mu + \delta_1)^2}{\left\| (q\beta_2 S(t) I_2(t))^2 \right\|_\infty + \left\| (\gamma E(t))^2 \right\|_\infty} |I_1(t)|^2 \right) \end{aligned}$$

under the condition

$$\frac{\left\| (p\beta_1 S(t))^2 \right\|_\infty + (\phi + \mu + \delta_1)^2}{\left\| (q\beta_2 S(t) I_2(t))^2 \right\|_\infty + \left\| (\gamma E(t))^2 \right\|_\infty} < 1, \text{ then} \quad (42)$$

$$|f_3(S, E, I_1, I_2)|^2 \leq k_3(1 + |I_1(t)|^2). \quad (43)$$

Finally for the function $f_4(S, E, I_1, I_2)$

$$\begin{aligned} |f_4(S, E, I_1, I_2)|^2 &= |\phi(1 - r_1) I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t)|^2, \quad (44) \\ &\leq 3 |\phi(1 - r_1) I_1(t)|^2 + 3 |(\mu + \delta_2) I_2(t)|^2 \\ &\quad + 3 |\varphi r_2 I_2(t)|^2, \end{aligned}$$

$$\begin{aligned}
 &\leq 3\phi^2(1-r_1)|I_1(t)|^2 + 3(\mu + \delta_2)^2 |I_2(t)|^2 \\
 &+ (\varphi r_2)^2 3 |I_2(t)|^2, \\
 &\leq 3\phi^2(1-r_1) \sup_{t \in [0, T_2]} |I_1(t)|^2 + 3(\mu + \delta_2)^2 |I_2(t)|^2 \\
 &+ (\varphi r_2)^2 3 |I_2(t)|^2, \\
 &\leq 3\phi^2(1-r_1) \|I_1^2(t)\|_\infty + 3(\mu + \delta_2)^2 |I_2(t)|^2 + (\varphi r_2)^2 3 |I_2(t)|^2, \\
 &\leq 3\phi^2(1-r_1) \|I_1^2(t)\|_\infty \left(1 + \frac{(\mu + \delta_2)^2 + (\varphi r_2)^2}{\phi^2(1-r_1) \|I_1^2(t)\|_\infty} |I_2(t)|^2 \right),
 \end{aligned}$$

under the condition

$$\frac{(\mu + \delta_2)^2 + (\varphi r_2)^2}{\phi^2(1-r_1) \|I_1^2(t)\|_\infty} < 1, \text{ then} \tag{45}$$

$$|f_4(S, E, I_1, I_2)|^2 \leq k_4(1 + |I_2(t)|^2). \tag{46}$$

Therefore the condition of linear growth is verified if

$$\max \left\{ \begin{aligned} &\frac{\|(\beta_1 I_1(t))^2\|_\infty + \|(\beta_2 I_2(t))^2\|_\infty + \mu^2}{\lambda^2}, \\ &\frac{(\mu + \gamma)^2}{\|(\beta_1 p_1 S(t) I_1(t))^2\|_\infty + \|(\beta_2 q_1 S(t) I_2(t))^2\|_\infty}, \\ &\frac{\|(p \beta_1 S(t))^2\|_\infty + (\phi + \mu + \delta_1)^2}{\|(\beta_1 S(t) I_1(t))^2\|_\infty + \|(\beta_2 S(t) I_2(t))^2\|_\infty}, \\ &\frac{(\mu + \delta_2)^2 + (\varphi r_2)^2}{\phi^2(1-r_1) \|I_1^2(t)\|_\infty} \end{aligned} \right\} < 1. \tag{47}$$

Now we have to verify Lipschitz condition for equations.

For the function $f_1(S, E, I_1, I_2)$,

$$\begin{aligned}
 |f_1(S, E, I_1, I_2) - f_1(S', E, I_1, I_2)| &\leq (\beta_1 I_1(t) + \beta_2 I_2(t) + \mu) |S - S'|, \\
 &\leq \bar{k}_1 |S - S'|.
 \end{aligned} \tag{48}$$

For the function $f_2(S, E, I_1, I_2)$,

$$\begin{aligned}
 |f_2(S, E, I_1, I_2) - f_2(S, E', I_1, I_2)| &\leq (\gamma + \mu) |E - E'|, \\
 &\leq \bar{k}_2 |E - E'|.
 \end{aligned} \tag{49}$$

For the function $f_3(S, E, I_1, I_2)$,

$$\begin{aligned} |f_3(S, E, I_1, I_2) - f_3(S, E, I'_1, I_2)| &\leq (p\beta_1 S(t) + (\phi + \mu + \delta_1)) |I_1 - I'_1|, \\ &\leq \bar{k}_3 |I_1 - I'_1|. \end{aligned} \quad (50)$$

Finally for the function $f_4(S, E, I_1, I_2)$,

$$\begin{aligned} |f_4(S, E, I_1, I_2) - f_4(S, E, I_1, I'_2)| &\leq (\mu + \delta_2 + \varphi r_2) |I_2 - I'_2|, \\ &\leq \bar{k}_4 |I_2 - I'_2|. \end{aligned} \quad (51)$$

We verified the Lipschitz condition which completes the proof.

Let us do proof for last part of piecewise equation. Here we consider for $\forall t \in [W_2, W]$. In the model we take for $S(t), E(t), I_1(t), I_2(t) \in [W_2, \tau_e]$, where τ_e shows explosion time. To prove the solution is global, one has to prove that such system solution is global, so we have to prove that $\tau_e = \infty$.

Now we consider $l_0 \in R_+$ is a positive constant such that $S(W_2), E(W_2), I_1(W_2), I_2(W_2)$ lies within $[\frac{1}{l_0}, l_0]$. We define a stopping time as

$$\tau_l = \left\{ t \in [W_2, \tau_e] : \frac{1}{l} \geq \min\{S(t), E(t), I_1(t), I_2(t)\} \text{ or } \max\{S(t), E(t), I_1(t)I_2(t)\} \geq l \right\}, \quad (52)$$

for each $l \geq l_0$. While as $l \rightarrow \infty$, τ_l is monotonically increasing. $\lim_{l \rightarrow \infty} \tau_l = \tau_\infty$ with $\tau_e \geq \tau_\infty$. $\forall t \geq 0$, if

we show that $\tau_\infty = 0$, then we can conclude that $\tau_e = \infty$ and $S(t), E(t), I_1(t), I_2(t) \in R_+^4$ is solution. So we have to prove that $\tau_e = \infty$.

If we have contradictory for the conclusion, then there exists $0 < W$ and $\varepsilon \in (0, 1)$ such that $P\{W \geq \tau_\infty\} > \varepsilon$. (53)

Now we define a function $H(X) : R_+^4 \rightarrow R_+$ in $H \in C^2$ such that

$$\begin{aligned} \bar{H}(X) = dH(X) &= \sum_{j=1}^4 \left(1 - \frac{1}{x_j}\right) dx_j + \sum_{j=1}^4 \sigma_j (x_j - 1) dB_j(t), \\ &= \sum_{j=1}^4 \left(1 - \frac{1}{x_j}\right) x'_j + \sum_{j=1}^4 \sigma_j (x_j - 1) dB_j(t), \end{aligned} \quad (54)$$

where

$$x_1 = S(t), \quad x_2 = E(t), \quad x_3 = I_1(t), \quad x_4 = I_2(t),$$

$$\sigma_j = (\sigma_1, \sigma_2, \sigma_3, \sigma_4), \quad (55)$$

$$B_j(t) = (B_1(t), B_2(t), B_3(t), B_4(t)).$$

For our model $\bar{H}(X)$ is obtained by following equality:

$$\begin{aligned} \bar{H}(X) = & \sum_{j=1}^4 \left(1 - \frac{1}{x_j}\right) x'_j = \left(1 - \frac{1}{S}\right) S' + \left(1 - \frac{1}{E}\right) E' \\ & + \left(1 - \frac{1}{I_1}\right) I'_1 + \left(1 - \frac{1}{I_2}\right) I'_2 + \sum_{j=1}^4 \frac{\sigma_j^2}{2}. \end{aligned} \tag{56}$$

$$\begin{aligned} \bar{H}(X) = & \lambda + \beta_1 I_1(t) + \beta_2 I_2(t) + \mu \\ & + \beta_1 p_1 S(t) I_1(t) + \beta_2 q_1 S(t) I_2(t) \\ & + (\mu + \gamma) + p \beta_1 S(t) I_1(t) \\ & + \beta_2 q S(t) I_2(t) + \gamma E(t) + (\phi + \mu + \delta_1) \\ & + \phi(1 - r_1) I_1(t) + (\mu + \delta_2) I_2(t) + \varphi r_2 I_2(t) \\ & - \left\{ \begin{aligned} & \beta_1 S(t) I_1(t) + \beta_2 S(t) I_2(t) + \mu S(t) + \frac{\lambda}{S} + (\mu + \gamma) E(t) \\ & + \frac{1}{E} \beta_1 p_1 S(t) I_1(t) + \frac{1}{E} \beta_2 q_1 S(t) I_2(t) + (\phi + \mu + \delta_1) I_1(t) \\ & + p \beta_1 S(t) + \frac{1}{I_1} \beta_2 q S(t) I_2(t) + \frac{1}{I_1} \gamma E(t) \\ & + (\mu + \delta_2) I_2(t) + \varphi r_2 I_2(t) + \frac{1}{I_2} \phi(1 - r_1) I_1(t) \end{aligned} \right. \\ & + \sum_{j=1}^4 \frac{\sigma_j^2}{2}, \\ & < \lambda + 3\mu + \gamma + \phi + \delta_1 = \bar{\theta} \end{aligned} \tag{57}$$

and

$$\bar{H}(X) = \bar{\theta} dt + \sum_{j=1}^4 \sigma_j (x_j - 1) dB_j(t). \tag{58}$$

By taking integration from 0 to $\tau_l \wedge W$, we have

$$\begin{aligned} E[\bar{H}(\tau_l \wedge X)] \leq & \bar{H}(X(W_2)) + E \left[\int_0^{\tau_l \wedge W} \bar{\theta} \right], \\ & \leq \bar{H}(X(W_2)) + \bar{\theta} W. \end{aligned} \tag{59}$$

Setting $\Pi_l = \{W > \tau_l\}$ for $l_1 \leq l$ and thus $P(\Pi_l) \geq \zeta$.

Notting that for $\forall \Omega \in \Pi_l$, there must exist at least one $X(\tau_l, w)$ which is equal to $\frac{1}{7}$ or l . Then $l - \log l - 1$ or $\frac{1}{7} + \log l - 1$ as result

$$\left(\frac{1}{7} + \log l - 1\right) \wedge E(l - \log l - 1) < \overline{H}(X(\tau_l)). \quad (60)$$

From above, we can write

$$\begin{aligned} \overline{H}(X(W_2)) + \overline{\theta} W &> E(1_{\Pi_l} \overline{H}(X(\tau_l))), \\ &\geq \sigma \left[(l - \log l - 1) \wedge \left(\frac{1}{7} + \log l - 1\right) \right]. \end{aligned} \quad (61)$$

Here 1_{Π_l} is the indicator function of Π . Thus $\lim_{l \rightarrow \infty}$ leads

$$\infty > \overline{H}(X(W_2)) + \overline{\theta} W = 0. \quad (62)$$

It is a contradiction. So under the conditions gived earlier $\tau_\infty = \infty$ which completes the proof.

4 Numerical Schemes for Model with Four Waves Patterns

In this section, we generate a numerical schemes for spread of infectious (specially for pandemic) disease with four patterns. These schemes will consist of three derivatives with randomness [1,12].

4.1 Case 1: Classical-Power Law-Exponential Decay Law-Randomness

In this case, we consider a version with four waves which have classical derivative starts from 0 to W_1 , the power law derivative start from W_1 to W_2 , the exponential decay law derivative start from W_2 to W_3 , and the last from W_3 to W . So a piecewise mathematical system that is defined as subsection can be given as

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C_{t_1} D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{CF}_{t_2} D_t^\alpha y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3}, & i = 1, 2, \dots, n \end{array} \right. \quad (63)$$

where σ_i are densities of randomness and B_i are the functions of noise.

4.2 Case 2: Classical-Power Law-Mittag-Leffler Law-Randomness

In this case, we consider a version with four waves which have classical derivative starts from 0 to W_1 , the power law derivative start from W_1 to W_2 , the Mittag-Leffler law derivative start from W_2 to W_3 , and the last from W_3 to W . So a piecewise mathematical system that is defined as subsection can be given as

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{ABC} D_t^\alpha y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. , \tag{64}$$

where σ_i are densities of randomness and B_i are the functions of noise.

4.3 Case 3: Classical-Power Law-Fractal-Fractional Power Law Derivative-Randomness

In this case, we consider a version with four waves which have classical derivative starts from 0 to W_1 , the power law derivative start from W_1 to W_2 , fractal-fractional power law derivative start from W_2 to W_3 , and the last from W_3 to W . So a piecewise mathematical system that is defined as subsection can be given as

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFP} D_t^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. , \tag{65}$$

where σ_i are densities of randomness and B_i are the functions of noise.

4.4 Case 4: Classical-Exponential Decay Law-Fractal-Fractional Exponential Decay Law Derivative-Randomness

In this case, we consider a version with four waves which have classical derivative starts from 0 to W_1 , the exponential decay law derivative start from W_1 to W_2 , fractal-fractional exponential decay law derivative start from W_2 to W_3 , and the last from W_3 to W . So a piecewise mathematical system that is defined as subsection can be given as

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^{CF}D_{t_1}^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFE}D_{t_2}^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \quad (66)$$

where σ_i are densities of randomness and B_i are the functions of noise.

4.5 Case 5: Classical-Mittag-Leffler Law-Fractal-Fractional Mittag-Leffler Law Derivative-Randomness

In this case, we consider a version with four waves which have classical derivative starts from 0 to W_1 , the Mittag Leffler law derivative start from W_1 to W_2 , fractal-fractional Mittag-Leffler law derivative start from W_2 to W_3 , and the last from W_3 to W . So a piecewise mathematical system that is defined as subsection can be given as

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^{ABC}D_{t_1}^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFM}D_{t_2}^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \quad (67)$$

where σ_i are densities of randomness and B_i are the functions of noise.

5 Numerical Schemes of Piecewise Epidemic Disease Models with Four Waves Patterns

In this section we assumed that those kind of epidemic models satisfy existence and uniqueness. So we can put numerical solutions for them. While putting solution results we use in all cases on the Lagrange polynomial interpolation. First we divide $[0, W]$ in four

$$0 \leq t_0 \leq t_1 \leq \dots \leq t_{n_1} = W_1 \leq t_{n_1+1} \leq t_{n_1+2} \leq \dots \leq t_{n_2} = W_2 \tag{68}$$

$$\leq t_{n_2+1} \leq t_{n_2+2} \leq \dots \leq t_{n_3} = W_3 \leq t_{n_3+1} \leq t_{n_3+2} \leq \dots \leq t_{n_3} = W.$$

5.1 Numerical Method for Case 1:

Let us consider the first case

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{CF} D_t^\alpha y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \tag{69}$$

The numerical solution can then be provided as

$$\left\{ \begin{array}{l} y_i^{n_1} = y_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} g(t_{k_1}, y(t_{k_1})) - g(t_{k_1-1}, y(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\ y_i^{n_2} = y_i(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_2=0}^{n_2} g(t_{k_2}, y(t_{k_2})) \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix} \\ \quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_2=0}^{n_2} g(t_{k_2-1}, y(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\ y_i^{n_3} = y_i(W_2) + \frac{1-\alpha}{M(\alpha)} \sum_{k_3=0}^{n_3} [g(t_{k_3}, y(t_{k_3})) - g(t_{k_3-1}, y(t_{k_3-1}))] \\ \quad + \frac{\alpha}{M(\alpha)} \sum_{k_3=0}^{n_3} \left\{ \frac{3\Delta t}{2} g(t_{k_3}, y(t_{k_3})) - g(t_{k_3-1}, y(t_{k_3-1})) \frac{\Delta t}{2} \right\}, \quad W_2 \leq t \leq W_3 \\ y_i^{n_4} = y_i(W_3) + \sum_{k_4=i+1}^{n_4} \left\{ \frac{3\Delta t}{2} g(t_{k_4}, y(t_{k_4})) - g(t_{k_4-1}, y(t_{k_4-1})) \frac{\Delta t}{2} \right\} \quad W_3 \leq t \leq W \\ \quad + \sum_{k_4=i+1}^{n_4} \left\{ \frac{\sigma}{2} (y(t_{k_4+1}) + y(t_{k_4})) (B(t_{k_4+1}) - B(t_{k_4})) \right\}. \end{array} \right. \tag{70}$$

5.2 Numerical Method for Case 2:

We deal with the following problem with second case

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq T_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C D_t^\alpha y_i = g(t, y), & \text{if } T_1 \leq t \leq T_2 \\ y_i(T_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{ABC} D_t^\alpha y_i = g(t, y), & \text{if } T_2 \leq t \leq T_3 \\ y_i(T_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } T_3 \leq t \leq T \\ y_i(T_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \quad (71)$$

The numerical solution for such problem is given by

$$\left\{ \begin{array}{l} y_i^{n1} = y_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} g(t_{k_1}, y(t_{k_1})) - g(t_{k_1-1}, y(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\ y_i^{n2} = y_i(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} g(t_{k_2}, y(t_{k_2})) \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix} \\ \quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} g(t_{k_2-1}, y(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\ y_i^{n3} = y_i(W_2) + \frac{1-\alpha}{AB(\alpha)} f(t_{k_3}, y(t_{k_3})) \\ \quad + \frac{\alpha (\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_3=0}^{n_3} g(t_{k_3}, y(t_{k_3})) \times \begin{bmatrix} (n_3 - k_3 + 1)^\alpha (n_3 - k_3 + 2 + \alpha) \\ -(n_3 - k_3)^\alpha (n_3 - k_3 + 2 + 2\alpha) \end{bmatrix} \\ \quad - \frac{\alpha (\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_3=0}^{n_3} g(t_{k_3-1}, y(t_{k_3-1})) \times \left[(n_3 - k_3 + 1)^{\alpha+1} - (n_3 - k_3)^\alpha (n_3 - k_3 + 1 + \alpha) \right], \quad W_2 \leq t \leq W_3 \\ y_i^{n4} = y_i(W_3) + \sum_{k_4=i+1}^{n_4} \left\{ \frac{3\Delta t}{2} g(t_{k_4}, y(t_{k_4})) - g(t_{k_4-1}, y(t_{k_4-1})) \frac{\Delta t}{2} \right\} \quad W_3 \leq t \leq W \\ \quad + \sum_{k_4=i+1}^{n_4} \left\{ \frac{\sigma}{2} (y(t_{k_4+1}) + y(t_{k_4})) (B(t_{k_4+1}) - B(t_{k_4})) \right\}. \end{array} \right. \quad (72)$$

5.3 Numerical Method for Case 3:

Now we deal with the following problem with third case

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^C D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFP} D_t^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \quad (73)$$

The numerical solution for such problem is given by

$$\left\{ \begin{array}{l} y_i^{n_1} = y_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} g(t_{k_1}, y(t_{k_1})) - g(t_{k_1-1}, y(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\ y_i^{n_2} = y_i(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_2=0}^{n_2} g(t_{k_2}, y(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix} \\ \quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_2=0}^{n_2} g(t_{k_2-1}, y(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\ y_i^{n_3} = y_i(W_2) + \frac{\beta (\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_3=0}^{n_3} t_{k_3}^{\beta-1} g(t_{k_3}, y(t_{k_3})) \times \begin{bmatrix} (n_3 - k_3 + 1)^\alpha (n_3 - k_3 + 2 + \alpha) \\ -(n_3 - k_3)^\alpha (n_3 - k_3 + 2 + 2\alpha) \end{bmatrix} \\ \quad + \frac{\beta (\Delta t)^\alpha}{\Gamma(\alpha + 2)} \sum_{k_3=0}^{n_3} t_{k_3-1}^{\beta-1} g(t_{k_3-1}, y(t_{k_3-1})) \times \begin{bmatrix} (n_3 - k_3 + 1)^{\alpha+1} \\ -(n_3 - k_3)^\alpha (n_3 - k_3 + 1 + \alpha) \end{bmatrix}, \quad W_2 \leq t \leq W_3 \\ y_i^{n_4} = y_i(W_3) + \sum_{k_4=i+1}^{n_4} \left\{ \begin{array}{l} \frac{3\Delta t}{2} g(t_{k_4}, y(t_{k_4})) \\ -g(t_{k_4-1}, y(t_{k_4-1})) \frac{\Delta t}{2} \end{array} \right\} \\ \quad + \sum_{k_4=i+1}^{n_4} \left\{ \frac{\sigma}{2} (y(t_{k_4+1}) + y(t_{k_4})) (B(t_{k_4+1}) - B(t_{k_4})) \right\}. \quad W_3 \leq t \leq W \end{array} \right. \quad (74)$$

5.4 Numerical Method for Case 4:

Here, we deal with the following problem with fourth case

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^{CF}D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFE}D_t^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1 i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W, \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \quad (75)$$

The numerical solution for such problem is given by

$$\left\{ \begin{array}{ll} y_i^{n_1} = y_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} g(t_{k_1}, y(t_{k_1})) - g(t_{k_1-1}, y(t_{k_1-1})) \frac{\Delta t}{2} \right\}, & 0 \leq t \leq W_1 \\ y_i^{n_2} = y_i(W_1) + \frac{1-\alpha}{M(\alpha)} \sum_{k_2=0}^{n_2} [g(t_{k_2}, y(t_{k_2})) - g(t_{k_2-1}, y(t_{k_2-1}))] & W_1 \leq t \leq W_2 \\ \quad + \frac{\alpha}{M(\alpha)} \sum_{k_2=0}^{n_2} \left\{ \frac{3\Delta t}{2} g(t_{k_2}, y(t_{k_2})) - g(t_{k_2-1}, y(t_{k_2-1})) \frac{\Delta t}{2} \right\}, \\ y_i^{n_3} = y_i(W_2) + \frac{1-\alpha}{M(\alpha)} \sum_{k_3=0}^{n_3} [\beta t_{k_3}^{\beta-1} g(t_{k_3}, y(t_{k_3})) - \beta t_{k_3-1}^{\beta-1} g(t_{k_3-1}, y(t_{k_3-1}))] & W_2 \leq t \leq W_3 \\ \quad + \frac{\alpha}{M(\alpha)} \sum_{k_3=0}^{n_3} \left\{ \frac{3\Delta t}{2} \beta t_{k_3}^{\beta-1} g(t_{k_3}, y(t_{k_3})) - \beta t_{k_3-1}^{\beta-1} g(t_{k_3-1}, y(t_{k_3-1})) \frac{\Delta t}{2} \right\}, \\ y_i^{n_4} = y_i(W_3) + \sum_{k_4=i+1}^{n_4} \left\{ \frac{3\Delta t}{2} g(t_{k_4}, y(t_{k_4})) - g(t_{k_4-1}, y(t_{k_4-1})) \frac{\Delta t}{2} \right\} & W_3 \leq t \leq W \\ \quad + \sum_{k_4=i+1}^{n_4} \left\{ \frac{\sigma}{2} (y(t_{k_4+1}) + y(t_{k_4})) (B(t_{k_4+1}) - B(t_{k_4})) \right\}. \end{array} \right. \quad (76)$$

6 Numerical Method for Case 5:

Finally, we give numerical method with the following problem with fifth case:

$$\left\{ \begin{array}{ll} \frac{dy_i}{dt} = g(t, y), & \text{if } 0 \leq t \leq W_1 \\ y_i(0) = y_{i,0}, & i = 1, 2, \dots, n \\ {}^{ABC}D_t^\alpha y_i = g(t, y), & \text{if } W_1 \leq t \leq W_2 \\ y_i(W_1) = y_{i,1}, & 0 < \alpha \leq 1, i = 1, 2, \dots, n \\ {}^{FFM}D_t^{\alpha, \beta} y_i = g(t, y), & \text{if } W_2 \leq t \leq W_3 \\ y_i(W_2) = y_{i,2}, & 0 < \alpha \leq 1 i = 1, 2, \dots, n \\ dy(t) = g(t, y)dt + \sigma_i y_i dB_i(t), & \text{if } W_3 \leq t \leq W \\ y_i(W_3) = y_{i,3} & i = 1, 2, \dots, n \end{array} \right. \tag{77}$$

The numerical solution for such problem is given by

$$\left\{ \begin{array}{l} y_i^{n_1} = y_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} g(t_{k_1}, y(t_{k_1})) - g(t_{k_1-1}, y(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\ y_i^{n_2} = y_i(W_1) + \frac{1-\alpha}{AB(\alpha)} g(t_{n_2}, y(t_{n_2})) \\ + \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} g(t_{k_2}, y(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix} \quad W_1 \leq t \leq W_2 \\ - \frac{\alpha(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} g(t_{k_2-1}, y(t_{k_2-1})) \times \left[(n_2 - k_2 + 1)^{\alpha+1} - (n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \right], \\ y_i^{n_3} = y_i(W_2) + \frac{1-\alpha}{AB(\alpha)} t_{n_3}^{\beta-1} g(t_{n_3}, y(t_{n_3})) \quad W_2 \leq t \leq W_3 \\ + \frac{\alpha\beta(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_3=0}^{n_3} t_{k_3}^{\beta-1} g(t_{k_3}, y(t_{k_3})) \times \begin{bmatrix} (n_3 - k_3 + 1)^\alpha (n_3 - k_3 + 2 + \alpha) \\ -(n_3 - k_3)^\alpha (n_3 - k_3 + 2 + 2\alpha) \end{bmatrix} \\ + \frac{\alpha\beta(\Delta t)^\alpha}{AB(\alpha)\Gamma(\alpha+2)} \sum_{k_3=0}^{n_3} t_{k_3-1}^{\beta-1} g(t_{k_3-1}, y(t_{k_3-1})) \times \begin{bmatrix} (n_3 - k_3 + 1)^{\alpha+1} \\ -(n_3 - k_3)^\alpha (n_3 - k_3 + 1 + \alpha) \end{bmatrix}, \\ y_i^{n_4} = y_i(W_3) + \sum_{k_4=i+1}^{n_4} \left\{ \frac{3\Delta t}{2} g(t_{k_4}, y(t_{k_4})) - g(t_{k_4-1}, y(t_{k_4-1})) \frac{\Delta t}{2} \right\} \quad W_3 \leq t \leq W \\ + \sum_{k_4=i+1}^{n_4} \left\{ \frac{\sigma}{2} (y(t_{k_4+1}) + y(t_{k_4})) (B(t_{k_4+1}) - B(t_{k_4})) \right\}, \end{array} \right. \tag{78}$$

6.1 Numerical Simulation for Stochastic-Deterministic Model of Tuberculosis

In this section, we give numerical simulation of the Tuberculosis epidemic system of fractional stochastic differential equations. We have made use of the model with the piecewise differential

operators and the numerical scheme where the Lagrange polynomial interpolation is used. While modelling with piecewise idea, the first part is classical, the second part is fractional and last part is stochastic. The numerical simulation is performed for different values of fractional orders. So the stochastic-deterministic piecewise tuberculosis model is given as

$$\begin{cases} \frac{dS(t)}{dt} = \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\ \frac{dE(t)}{dt} = \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\ \frac{dI_1(t)}{dt} = p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\ \frac{dI_2(t)}{dt} = \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t), \\ S(0) = S_0, \quad E(0) = E_0, \quad I_1(0) = I_{10}, \quad I_2(0) = I_{20}, \end{cases} \quad \text{if } 0 \leq t \leq W_1 \quad (79)$$

$$\begin{cases} {}^C D_t^\alpha S(t) = \lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t), \\ {}^C D_t^\alpha E(t) = \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) - (\mu + \gamma) E(t), \\ {}^C D_t^\alpha I_1(t) = p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) - (\phi + \mu + \delta_1) I_1(t), \\ {}^C D_t^\alpha I_2(t) = \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) - \varphi r_2 I_2(t), \\ S(W_1) = S_1, \quad E(W_1) = E_1, \quad I_1(W_1) = I_{11}, \quad I_2(W_1) = I_{21}, \end{cases} \quad \text{if } W_1 \leq t \leq W_2 \quad (80)$$

$$\begin{cases} dS(t) = [\lambda - \beta_1 S(t)I_1(t) - \beta_2 S(t)I_2(t) - \mu S(t)] dt + \sigma_1 S dB_1(t), \\ dE(t) = \begin{bmatrix} \beta_1 p_1 S(t)I_1(t) - \beta_2 q_1 S(t)I_2(t) \\ -(\mu + \gamma) E(t) \end{bmatrix} dt + \sigma_2 E dB_2(t), \\ dI_1(t) = \begin{bmatrix} p\beta_1 S(t)I_1(t) + q\beta_2 S(t)I_2(t) + \gamma E(t) \\ -(\phi + \mu + \delta_1) I_1(t) \end{bmatrix} dt + \sigma_3 I_1 dB_3(t), \\ dI_2(t) = \begin{bmatrix} \phi(1 - r_1)I_1(t) - (\mu + \delta_2) I_2(t) \\ -\varphi r_2 I_2(t) \end{bmatrix} dt + \sigma_4 I_2 dB_4(t), \\ S(W_2) = S_2, \quad E(W_2) = E_2, \quad I_1(W_2) = I_{12}, \quad I_2(W_2) = I_{22}, \end{cases} \quad \text{if } W_2 \leq t \leq W. \quad (81)$$

For simplicity we consider right side of system as

$$\begin{cases} \dot{S} = f_1(S, E, I_1, I_2), \\ \dot{E} = f_2(S, E, I_1, I_2), \\ \dot{I}_1 = f_3(S, E, I_1, I_2), \\ \dot{I}_2 = f_4(S, E, I_1, I_2), \end{cases} \quad (82)$$

Using the numerical scheme presented in this paper with piecewise derivative, the numerical solution of the stochastic-deterministic tuberculosis model is given as follows:

$$\left\{ \begin{aligned}
 S_i^{n1} &= S_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} f_1(t_{k_1}, S(t_{k_1})) - f_1(t_{k_1-1}, S(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\
 S_i^{n2} &= S_i(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_1(t_{k_2}, S(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\
 &\quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_1(t_{k_2-1}, S(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \\
 S_i^{n3} &= S_i(W_2) + \sum_{k_3=i+1}^{n_3} \left\{ \begin{array}{l} \frac{3\Delta t}{2} f_1(t_{k_3}, S(t_{k_3})) \\ -f_1(t_{k_3-1}, S(t_{k_3-1})) \frac{\Delta t}{2} \end{array} \right\} \quad W_2 \leq t \leq W \\
 &\quad + \sum_{k_3=i+1}^{n_3} \left\{ \frac{\sigma}{2} (S(t_{k_3+1}) + S(t_{k_3})) (B(t_{k_3+1}) - B(t_{k_3})) \right\}, \\
 \\
 E_i^{n1} &= E_i(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} f_2(t_{k_1}, E(t_{k_1})) - f_2(t_{k_1-1}, E(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\
 E_i^{n2} &= E_i(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_2(t_{k_2}, E(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix}, \quad T_1 \leq t \leq T_2 \\
 &\quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_2(t_{k_2-1}, E(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix} \\
 E_i^{n3} &= E_i(W_2) + \sum_{k_3=i+1}^{n_3} \left\{ \begin{array}{l} \frac{3\Delta t}{2} f_2(t_{k_3}, E(t_{k_3})) \\ -f_2(t_{k_3-1}, E(t_{k_3-1})) \frac{\Delta t}{2} \end{array} \right\} \quad W_2 \leq t \leq W. \\
 &\quad + \sum_{k_3=i+1}^{n_3} \left\{ \frac{\sigma}{2} (E(t_{k_3+1}) + E(t_{k_3})) (B(t_{k_3+1}) - B(t_{k_3})) \right\} \\
 \\
 I_{1i}^{n1} &= I_{1i}(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} f_3(t_{k_1}, I_1(t_{k_1})) - f_3(t_{k_1-1}, I_1(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1, \\
 I_{1i}^{n2} &= I_{1i}(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_3(t_{k_2}, I_1(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\
 &\quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_3(t_{k_2-1}, I_1(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \\
 I_{1i}^{n3} &= I_{1i}(W_3) + \sum_{k_3=i+1}^{n_3} \left\{ \begin{array}{l} \frac{3\Delta t}{2} f_3(t_{k_3}, I_1(t_{k_3})) \\ -f_3(t_{k_3-1}, I_1(t_{k_3-1})) \frac{\Delta t}{2} \end{array} \right\} \quad W_2 \leq t \leq W. \\
 &\quad + \sum_{k_3=i+1}^{n_3} \left\{ \frac{\sigma}{2} (I_1(t_{k_3+1}) + I_1(t_{k_3})) (B(t_{k_3+1}) - B(t_{k_3})) \right\},
 \end{aligned} \right. \tag{83}$$

$$\left\{ \begin{aligned}
 I_{2i}^{n_1} &= I_{2i}(0) + \sum_{k_1=i+1}^{n_1} \left\{ \frac{3\Delta t}{2} f_4(t_{k_1}, I_2(t_{k_1})) - f_4(t_{k_1-1}, I_2(t_{k_1-1})) \frac{\Delta t}{2} \right\}, \quad 0 \leq t \leq W_1 \\
 I_{2i}^{n_2} &= I_{2i}(W_1) + \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_4(t_{k_2}, I_2(t_{k_2})) \times \begin{bmatrix} (n_2 - k_2 + 1)^\alpha (n_2 - k_2 + 2 + \alpha) \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 2 + 2\alpha) \end{bmatrix} \\
 &\quad - \frac{(\Delta t)^\alpha}{\Gamma(\alpha+2)} \sum_{k_2=0}^{n_2} f_4(t_{k_2-1}, I_2(t_{k_2-1})) \times \begin{bmatrix} (n_2 - k_2 + 1)^{\alpha+1} \\ -(n_2 - k_2)^\alpha (n_2 - k_2 + 1 + \alpha) \end{bmatrix}, \quad W_1 \leq t \leq W_2 \\
 I_{2i}^{n_3} &= I_{2i}(W_2) + \sum_{k_3=i+1}^{n_3} \left\{ \begin{aligned}
 &\frac{3\Delta t}{2} f_4(t_{k_3}, I_2(t_{k_3})) \\
 &- f_4(t_{k_3-1}, I_2(t_{k_3-1})) \frac{\Delta t}{2} \end{aligned} \right\} \quad W_2 \leq t \leq W \\
 &+ \sum_{k_3=i+1}^{n_3} \left\{ \frac{\sigma}{2} (I_2(t_{k_3+1}) + I_2(t_{k_3})) (B(t_{k_3+1}) - B(t_{k_3})) \right\}.
 \end{aligned} \right.$$

7 Numerical Simulations

In this section, we will deal with numerical simulation of the Tuberculosis epidemic system of fractional stochastic differential equations. in order to demonstrate that the proposed method is effective and accurate. We have made use of the model with the piecewise differential operators and the numerical scheme where the Lagrange polynomial interpolation is used. In the numerical scheme, the first part is classical, the second part is fractional and last part is stochastic. We also present the results obtained from the fractional stochastic model, the numerical simulations are shown in Fig. 1 for alpha = 1, Fig. 2 for alpha = 0.5, Fig. 3 for alpha = 0.6 and finally Fig. 4 for alpha = 0.9 with density of randomness given by sigma1 = 0.01, sigma2 = 0.015, sigma3 = 0.012, sigma4 = 0.010. And with same alpha values but different density of randomness given by sigma1 = 0.1, sigma2 = 0.2, sigma3 = 0.3, sigma4 = 0.4 we put Figs. 5–8. Also figures including the initial conditions as

$$S(1) = 180, \quad E(1) = 130, \quad I_1(1) = 160, \quad I_2(1) = 140.$$




Figure 1: Numerical simulation for alpha = 1




Figure 2: Numerical simulation for alpha = 0.5




Figure 3: Numerical simulation for $\alpha = 0.6$




Figure 4: Numerical simulation for $\alpha = 0.9$




Figure 5: Numerical simulation for $\alpha = 1$




Figure 6: Numerical simulation for $\alpha = 0.5$




Figure 7: Numerical simulation for $\alpha = 0.6$




Figure 8: Numerical simulation for $\alpha = 0.9$

8 Conclusion

The spread of tuberculosis within human settlements and has infected and killed millions of humans in the last 200 years. While researchers from all backgrounds have put their efforts together to combat this virus and try to stop its spread, several studies have been performed; however, the virus is still spreading so far. Mathematical models are used to predict the future development of a given real-world problem. While several techniques and models have been proposed, they have not predicted piecewise behaviors of the spread. In this work, we attempted to present a model with piecewise patterns.

Funding Statement: The authors received no specific funding for this study.

Conflicts of Interest: The authors declare that they have no conflicts of interest to report regarding the present study.

References

1. Atangana, A., Araz, S. S. (2021). Modeling and forecasting the spread of Covid-19 with stochastic and deterministic approaches: Africa and Europe. *Advances in Difference Equations*, 2021(1), 1–107. DOI 10.1186/s13662-021-03213-2.
2. Atangana, A. (2021). Mathematical model of survival of fractional calculus, critics and their impact: How singular is our world?. <https://hal.archives-ouvertes.fr/hal-03182789/document>.
3. Nematollahi, M. H., Vatankhah, R., Sharifi, M. (2020). Nonlinear adaptive control of tuberculosis with consideration of the risk of endogenous reactivation and exogenous reinfection. *Journal of Theoretical Biology*, 486, 110081. DOI 10.1016/j.jtbi.2019.110081.
4. Ogundile, O., Edeki, S., Adewale, S. (2018). A modelling technique for controlling the spread of tuberculosis. *International Journal of Biology and Biomedical Engineering*, 12, 131–136.
5. Atangana, A., Baleanu, D. (2016). New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model. *Thermal Science*, 20(2), 763. DOI 10.2298/TSCI160111018A.
6. Caputo, M., Fabrizio, M. (2017). On the notion of fractional derivative and applications to the hysteresis phenomena. *Meccanica*, 52(13), 3043–3052. DOI 10.1007/s11012-017-0652-y.
7. Caputo, M. (1967). Linear models of dissipation whose Q is almost frequency independent – II. *Geophysical Journal International*, 13(5), 529–539. DOI 10.1111/j.1365-246X.1967.tb02303.x.
8. Dokuyucu, M. A. (2020). Caputo and atangana-baleanu-caputo fractional derivative applied to garden equation. *Turkish Journal of Science*, 5(1), 1–7.
9. Dokuyucu, M. A., Celik, E. (2021). Analyzing a novel coronavirus model (Covid-19) in the sense of caputo-fabrizio fractional operator. *Applied and Computational Mathematics*, 20(1), 49–69.
10. Ilknur, K., AKçetin, E., Yaprakdal, P. (2020). Numerical approximation for the spread of siqr model with caputo fractional order derivative. *Turkish Journal of Science*, 5(2), 124–139.
11. Dokuyucu, M. A. (2020). Analysis of a fractional plant-nectar-pollinator model with the exponential kernel. *Eastern Anatolian Journal of Science*, 6(1), 11–20.
12. Atangana, A., Araz, S. S. (2021). New concept in calculus: Piecewise differential and integral operators. *Chaos, Solitons & Fractals*, 145, 110638. DOI 10.1016/j.chaos.2020.110638.
13. Qomariyah, N., Herdiana, R., Utomo, R., Permatasari, A. (2021). Stability analysis of a tuberculosis epidemic model with nonlinear incidence rate and treatment effects. *Journal of Physics: Conference Series*. 1943(1). 012118.
14. https://en.wikipedia.org/wiki/second_derivative.